

INERTIAL MANIFOLDS ON SQUEEZED DOMAINS

MARTINO PRIZZI
AND
KRZYSZTOF P. RYBAKOWSKI

ABSTRACT. Let Ω be an arbitrary smooth bounded domain in $\mathbb{R}^2$ and $\epsilon > 0$ be arbitrary. Squeeze Ω by the factor ϵ in the y -direction to obtain the squeezed domain $\Omega_\epsilon = \{(x, \epsilon y) \mid (x, y) \in \Omega\}$. In this paper we study the family of reaction-diffusion equations

$$(E_\epsilon) \quad \begin{aligned} u_t &= \Delta u + f(u), & t > 0, (x, y) \in \Omega_\epsilon \\ \partial_{\nu_\epsilon} u &= 0, & t > 0, (x, y) \in \partial\Omega_\epsilon, \end{aligned}$$

where f is a dissipative nonlinearity of polynomial growth. In a previous paper we showed that, as $\epsilon \rightarrow 0$, the equations (E_ϵ) have a limiting equation which is an abstract semilinear parabolic equation defined on a closed linear subspace of $H^1(\Omega)$. We also proved that the family $\mathcal{A}_\epsilon$ of the corresponding attractors is upper semicontinuous at $\epsilon = 0$. In this paper we prove that, if Ω satisfies some natural assumptions, then there is a family $\mathcal{M}_\epsilon$ of inertial C^1 -manifolds for (E_ϵ) of some fixed finite dimension ν . Moreover, as $\epsilon \rightarrow 0$, the flow on $\mathcal{M}_\epsilon$ converges in the C^1 -sense to the limit flow on $\mathcal{M}_0$.

1. INTRODUCTION

Let Ω be an arbitrary smooth bounded domain in $\mathbb{R}^2$ and $\epsilon > 0$ be arbitrary. Write (x, y) for a generic point of $\mathbb{R}^2$. Given $\epsilon > 0$ squeeze Ω by the factor ϵ in the y -direction to obtain the squeezed domain Ω_ϵ . More precisely, let $T_\epsilon: \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (x, \epsilon y)$ and $\Omega_\epsilon := T_\epsilon(\Omega)$.

Consider the following reaction-diffusion equation on Ω_ϵ :

$$(1_\epsilon) \quad \begin{aligned} u_t &= \Delta u + f(u), & t > 0, (x, y) \in \Omega_\epsilon \\ \partial_{\nu_\epsilon} u &= 0, & t > 0, (x, y) \in \partial\Omega_\epsilon. \end{aligned}$$

Here, ν_ϵ is the exterior normal vector field on $\partial\Omega_\epsilon$ and $f: \mathbb{R} \rightarrow \mathbb{R}$ is a C^1 -nonlinearity of polynomial growth such that $\limsup_{|s| \rightarrow \infty} f(s)/s \leq -\zeta$ for some $\zeta > 0$. These

hypotheses imply that (1_ϵ) generates a semiflow $\tilde{\pi}_\epsilon = \tilde{\pi}_{\epsilon,f}$ on $H^1(\Omega_\epsilon)$ which has a global attractor $\tilde{\mathcal{A}}_\epsilon = \tilde{\mathcal{A}}_{\epsilon,f}$.

As $\epsilon \rightarrow 0$ the thin domain Ω_ϵ degenerates to a one-dimensional interval.

One may ask what happens in the limit to the family $(\tilde{\pi}_\epsilon)_{\epsilon>0}$ of semiflows and to the family $(\tilde{\mathcal{A}}_\epsilon)_{\epsilon>0}$ of attractors. Is there a limit semiflow and a corresponding limit attractor?

This problem was first considered by Hale and Raugel in [7] for the case when the domain Ω is the *ordinate set* of a smooth positive function g defined on an interval $[a, b]$, i.e.

$$\Omega = \{ (x, y) \mid a < x < b \text{ and } 0 < y < g(x) \}.$$

The authors prove that, in this case, there exists a limit semiflow $\tilde{\pi}_0$, which is defined by the one-dimensional boundary value problem

$$(1_0) \quad \begin{aligned} u_t &= (1/g)(gu_x)_x + f(u), & t > 0, x \in]a, b[\\ u_x &= 0, & t > 0, x = a, b. \end{aligned}$$

Moreover, $\tilde{\pi}_0$ has a global attractor $\tilde{\mathcal{A}}_0$ and, in some sense, the family $(\tilde{\mathcal{A}}_\epsilon)_{\epsilon \geq 0}$ is upper-semicontinuous at $\epsilon = 0$.

Hale and Raugel also prove that one can modify the nonlinearity f in such a way that each modified semiflow $\tilde{\pi}'_\epsilon$ possesses an invariant C^1 -manifold $\tilde{\mathcal{M}}_\epsilon$ of some fixed dimension ν which includes the attractor $\tilde{\mathcal{A}}_\epsilon$ of the original semiflow $\tilde{\pi}_\epsilon$. The semiflows $\tilde{\pi}_\epsilon$ and $\tilde{\pi}'_\epsilon$ coincide on the attractor $\tilde{\mathcal{A}}_\epsilon$.

Moreover, as $\epsilon \rightarrow 0$, the reduced flow on $\tilde{\mathcal{M}}_\epsilon$ C^1 -converges to the reduced flow on $\tilde{\mathcal{M}}_0$.

If the domain Ω is not the ordinate set of some function (e.g. if Ω has holes or different horizontal branches) then (1_0) can no longer be a limiting equation for (1_ϵ) . Nevertheless, as it was proved in [15] the family $\tilde{\pi}_\epsilon$ still has a limit semiflow. Moreover, there exists a limit global attractor and the upper-semicontinuity result continues to hold.

In order to describe the main results of [15] we first transfer the family (1_ϵ) to boundary value problems on the fixed domain Ω . More explicitly, we use the linear isomorphism $\Phi_\epsilon: H^1(\Omega_\epsilon) \rightarrow H^1(\Omega)$, $u \mapsto u \circ T_\epsilon$, to transform problem (1_ϵ) to the equivalent problem

$$(2_\epsilon) \quad \begin{aligned} u_t &= u_{xx} + \frac{1}{\epsilon^2} u_{yy} + f(u), & t > 0, (x, y) \in \Omega \\ u_x \nu_1 + \frac{1}{\epsilon^2} u_y \nu_2 &= 0, & t > 0, (x, y) \in \partial\Omega. \end{aligned}$$

on Ω . Here, $\nu = (\nu_1, \nu_2)$ is the exterior normal vector field on $\partial\Omega$.

Note that equation (2_ϵ) can be written in the abstract form

$$\dot{u} + A_\epsilon u = \hat{f}(u)$$

where $\hat{f}: H^1(\Omega) \rightarrow L^2(\Omega)$ is the Nemitski operator generated by the function f , and A_ϵ is the linear operator defined by

$$A_\epsilon u = -u_{xx} - \frac{1}{\epsilon^2} u_{yy} \in L^2(\Omega) \quad \text{for } u \in H^2(\Omega) \text{ with } u_x \nu_1 + \frac{1}{\epsilon^2} u_y \nu_2 = 0 \text{ on } \partial\Omega.$$

Equation (2_ϵ) defines a semiflow π_ϵ on $H^1(\Omega)$ which is equivalent to $\tilde{\pi}_\epsilon$ and has the global attractor $\mathcal{A}_\epsilon := \Phi_\epsilon(\tilde{\mathcal{A}}_\epsilon)$, consisting of the orbits of all full bounded solutions of (2_ϵ) .

The operator A_ϵ is, in the usual way, induced by the following bilinear form

$$a_\epsilon(u, v) := \int_{\Omega} (u_x v_x + \frac{1}{\epsilon^2} u_y v_y) \, dx \, dy, \quad u, v \in H^1(\Omega).$$

Notice that, for every fixed $\epsilon > 0$ and $u \in H^1(\Omega)$, the formula

$$|u|_\epsilon = \left(a_\epsilon(u, u) + |u|_{L^2(\Omega)}^2 \right)^{1/2}$$

defines a norm on $H^1(\Omega)$ which is equivalent to $|\cdot|_{H^1(\Omega)}$. However, $|u|_\epsilon \rightarrow \infty$ as $\epsilon \rightarrow 0^+$ whenever $u_y \neq 0$ in $L^2(\Omega)$.

In fact, we see that for $u \in H^1(\Omega)$

$$\lim_{\epsilon \rightarrow 0^+} a_\epsilon(u, u) = \begin{cases} \int_{\Omega} u_x^2 \, dx \, dy, & \text{if } u_y = 0 \\ +\infty, & \text{otherwise.} \end{cases}$$

Thus the family $a_\epsilon(u, u)$, $\epsilon > 0$, of real numbers has a finite limit (as $\epsilon \rightarrow 0$) if and only if $u \in H_s^1(\Omega)$, where we define

$$H_s^1(\Omega) := \{ u \in H^1(\Omega) \mid u_y = 0 \}.$$

This is a closed linear subspace of $H^1(\Omega)$.

The corresponding limit bilinear form is given by the formula:

$$a_0(u, v) := \int_{\Omega} u_x v_x \, dx \, dy, \quad u, v \in H_s^1(\Omega).$$

The form a_0 uniquely determines a densely defined selfadjoint linear operator

$$A_0: D(A_0) \subset H_s^1(\Omega) \rightarrow L_s^2(\Omega)$$

by the usual formula

$$a_0(u, v) = \langle A_0 u, v \rangle_{L_s^2(\Omega)}, \quad \text{for } u \in D(A_0) \text{ and } v \in H_s^1(\Omega).$$

Here, $L_s^2(\Omega)$ is the closure of $H_s^1(\Omega)$ in the L^2 -norm, so $L_s^2(\Omega)$ is a closed linear subspace of $L^2(\Omega)$.

It follows that the Nemitski operator $\hat{f}$ maps the space $H_s^1(\Omega)$ into $L_s^2(\Omega)$. Consequently the abstract parabolic equation

$$(2_0) \quad \dot{u} = -A_0 u + \hat{f}(u)$$

defines a semiflow π_0 on the space $H_s^1(\Omega)$. This is the limit semiflow of the family π_ϵ . In fact, the following results are proved in [15]:

Theorem A. *Let $(\epsilon_n)_{n \in \mathbb{N}}$ be an arbitrary sequence of positive numbers convergent to zero and $(u_n)_{n \in \mathbb{N}}$ be a sequence in $L^2(\Omega)$ converging in the norm of $L^2(\Omega)$ to some $u_0 \in L^2_s(\Omega)$. Moreover, let $(t_n)_{n \in \mathbb{N}}$ be an arbitrary sequence of positive numbers converging to some positive number t_0 .*

Then

$$|e^{-t_n A_{\epsilon_n}} u_n - e^{-t_0 A_0} u_0|_{\epsilon_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

If, in addition, $u_n \in H^1(\Omega)$ for every $n \in \mathbb{N}$ and if $u_0 \in H^1_s(\Omega)$, then

$$|u_n \pi_{\epsilon_n} t_n - u_0 \pi_0 t_0|_{\epsilon_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

The limit semiflow π_0 possesses a global attractor $\mathcal{A}_0$. The upper-semicontinuity result alluded to above reads as follows:

Theorem B. *The family of attractors $(\mathcal{A}_\epsilon)_{\epsilon \in [0,1]}$ is upper-semicontinuous at $\epsilon = 0$ with respect to the family of norms $|\cdot|_\epsilon$.*

This means that

$$\lim_{\epsilon \rightarrow 0^+} \sup_{u \in \mathcal{A}_\epsilon} \inf_{v \in \mathcal{A}_0} |u - v|_\epsilon = 0.$$

In particular, there exists an $\epsilon_1 > 0$ and an open bounded set U in $H^1(\Omega)$ including all the attractors $\mathcal{A}_\epsilon$, $\epsilon \in [0, \epsilon_1]$.

The definition of the linear operator A_0 , as given above, is not very explicit. However, as it is shown in [15], there is a large class of the so-called *nicely decomposed* domains (see Figure 1) on which A_0 can be characterized as a system of one-dimensional second order linear differential operators, coupled to each other by certain compatibility and Kirchhoff type balance conditions. In this case, the abstract limit equation (2₀) is equivalent to a parabolic equation on a finite graph (see Figure 2).

Let $(\lambda_\nu)_{\nu \in \mathbb{N}}$ be the nondecreasing sequence of the eigenvalues of the limit operator A_0 (each of the eigenvalues being repeated according to its multiplicity). The first main result of the present paper says that given a nicely decomposed domain Ω satisfying some natural additional condition it follows that the sequence $(\lambda_\nu)_{\nu \in \mathbb{N}}$ satisfies the following gap condition:

$$(1.1) \quad \limsup_{\nu \rightarrow \infty} \frac{\lambda_{\nu+1} - \lambda_\nu}{\lambda_\nu^{1/2}} > 0.$$

(Cf. Theorem 4.8.)

The hypotheses of Theorem 4.8 are, in particular, satisfied in the ordinate set case considered in [7].

Now the second main theorem in this paper is the Inertial Manifold Theorem 5.2. This theorem says that under hypothesis (1.1) there is an $\epsilon_0 > 0$, $\epsilon_0 \leq \epsilon_1$, and there exists a family $\mathcal{M}_\epsilon$, $0 \leq \epsilon \leq \epsilon_0$ of (inertial) C^1 -manifolds of some finite dimension ν such that, whenever $0 \leq \epsilon \leq \epsilon_0$, then $\mathcal{A}_\epsilon \subset \mathcal{M}_\epsilon$ and the manifold $\mathcal{M}_\epsilon$ is locally invariant relative to the semiflow π_ϵ on the neighborhood U of the attractor $\mathcal{A}_\epsilon$.

Furthermore, as $\epsilon \rightarrow 0$, the reduced flow on the manifold $\mathcal{M}_\epsilon$ converges in the C^1 -sense to the reduced flow on $\mathcal{M}_0$.

In particular, our inertial manifold theorem contains, as a special case, the inertial manifold theorem of Hale and Raugel and it even improves the latter. In fact, for each $\epsilon > 0$ small enough, our inertial manifold $\mathcal{M}_\epsilon$ is (globally) invariant with respect to some modified semiflow π_ϵ'' coinciding with the original semiflow π_ϵ on the neighborhood U of the attractor $\mathcal{A}_\epsilon$. Thus, close to the attractor, $\mathcal{M}_\epsilon$ is a locally invariant manifold for the ‘true’ semiflow π_ϵ . On the other hand, the modified semiflow considered in [7] is, in general, different from the original semiflow on every H^1 -neighborhood of the attractor.

In order to prove their result, Hale and Raugel first develop H^2 -estimates for the attractors, which in turn imply the corresponding L^∞ -estimates. Then they modify the nonlinearity $f: \mathbb{R} \rightarrow \mathbb{R}$ so as to obtain a bounded nonlinearity $\tilde{f}: \mathbb{R} \rightarrow \mathbb{R}$ which induces a globally Lipschitzian Nemitski operator from $L^2(\Omega)$ to itself. Then they prove the existence of inertial manifolds in the space $L^2(\Omega)$ using the ‘cone-squeezing’ technique. This modification of the *function* f is the main reason why the modified semiflow $\tilde{\pi}'_\epsilon = \tilde{\pi}_{\epsilon, \tilde{f}}$ coincides with the original semiflow $\tilde{\pi}_\epsilon = \tilde{\pi}_{\epsilon, f}$ only on an H^2 -neighborhood the attractor.

In the present paper we apply the well-known technique for proving existence of invariant manifolds based on the contraction principle in an appropriate space of functions of exponential growth. In some cases, this method can be applied in a direct way to obtain existence of inertial manifolds (see e.g. [4], cf. also [18]). Unfortunately, such a direct application of this method to the present case does not work, since the gap condition (1.1), which is the best one possible, does not produce gaps which are ‘big enough’ to counterbalance the Lipschitz constant produced by the given Nemitski operator. As a consequence, the nonlinear operator Γ_ϵ used to define the inertial manifold $\mathcal{M}_\epsilon$ is **not** a contraction with respect to the natural norm $|u|_\epsilon$.

At this point we use a beautiful idea due to Brunovský and Tereščák [3]. This idea consists in replacing the norm $|u|_\epsilon$ by the equivalent norm

$$\|u\|_\epsilon = L|u|_{L^2} + l|u|_\epsilon.$$

This new norm depends on two positive constants L and l and one tries to determine these constants in such a way that the operator Γ_ϵ is a contraction with respect to the new norm. That such a choice is possible in our case follows by an application of the Gagliardo-Nirenberg inequality. Moreover, instead of modifying the nonlinearity f we only need to modify the corresponding Nemitski operator $\hat{f}: H^1(\Omega) \rightarrow L^2(\Omega)$, so as to obtain a globally Lipschitzian nonlinear operator $g: H^1(\Omega) \rightarrow L^2(\Omega)$ which coincides with $\hat{f}$ on U . This is the reason why the modified semiflow coincides on U with the original semiflow.

The reader is referred to the Reference section for some additional papers on thin domain problems. More extensive bibliography is contained in the survey paper [17] by G. Raugel.

In this paper we use standard notation, writing $\mathbb{R}$ and $\mathbb{N}$ to denote the set of reals and positive integers, respectively. We also identify $\mathbb{R} \times \mathbb{R}$ with $\mathbb{R}^2$, writing (x, y) for a generic point of $\mathbb{R}^2$. Finally, we denote by $\mathcal{L}^1$ and by $\mathcal{L}^2$ the Lebesgue measure in $\mathbb{R}$ and $\mathbb{R}^2$ respectively.

2. PRELIMINARIES

In this section we recall a few definitions and notations from [15]. We also discuss the existence and differentiability of Nemitski operators generated by real function.

Let Ω be a bounded domain in $\mathbb{R}^2$ with Lipschitz boundary.

For $\epsilon > 0$ define the symmetric bilinear form

$$a_\epsilon: H^1(\Omega) \times H^1(\Omega) \rightarrow \mathbb{R}$$

by

$$a_\epsilon(u, v) := \int_{\Omega} \left(\nabla_x u \cdot \nabla_x v + \frac{1}{\epsilon^2} \nabla_y u \cdot \nabla_y v \right) dx dy.$$

Let b be the restriction of the scalar product $\langle \cdot, \cdot \rangle_{L^2(\Omega)}$ to $H^1(\Omega) \times H^1(\Omega)$.

Since Ω has Lipschitz boundary, $H^1(\Omega)$ is dense in $L^2(\Omega)$ and the inclusion operator $H^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact. Therefore the pair (a_ϵ, b) defines a self-adjoint linear operator $A_\epsilon: D(A_\epsilon) \subset L^2(\Omega) \rightarrow L^2(\Omega)$ with domain dense in $L^2(\Omega)$.

Moreover, there are a nondecreasing sequence of eigenvalues $(\lambda_{\epsilon, j})_{j \in \mathbb{N}}$ and a corresponding L^2 -complete and L^2 -orthonormal system $(w_{\epsilon, j})_{j \in \mathbb{N}}$ of eigenvectors of the pair (a_ϵ, b) (equivalently, of A_ϵ).

On $H^1(\Omega)$ define the norm

$$|u|_\epsilon := \left(a_\epsilon(u, u) + |u|_{L^2(\Omega)}^2 \right)^{1/2}.$$

This norm is equivalent to $|\cdot|_{H^1(\Omega)}$ for every fixed $\epsilon > 0$, but $|u|_\epsilon \rightarrow \infty$ as $\epsilon \rightarrow 0^+$ whenever $\nabla_y u \neq 0$ in $L^2(\Omega)$.

Define the space

$$H_s^1(\Omega) = \{ u \in H^1(\Omega) \mid \nabla_y u = 0 \}.$$

Note that $H_s^1(\Omega)$ is a closed linear subspace of $H^1(\Omega)$.

Let $L_s^2(\Omega)$ to be the closure of the set $H_s^1(\Omega)$ in $L^2(\Omega)$. It follows that $L_s^2(\Omega)$ is a Hilbert space under the scalar product of $L^2(\Omega)$.

Note that $|u|_\epsilon \equiv |u|_{H^1(\Omega)}$ for $u \in H_s^1(\Omega)$.

Let $a_0: H_s^1(\Omega) \times H_s^1(\Omega) \rightarrow \mathbb{R}$ be the "limit" bilinear form defined by

$$a_0(u, v) := \int_{\Omega} \nabla u \cdot \nabla v dx dy = \int_{\Omega} \nabla_x u \cdot \nabla_x v dx dy.$$

Let b_0 be the restriction of the scalar product $\langle \cdot, \cdot \rangle_{L^2(\Omega)}$ to $H_s^1(\Omega) \times H_s^1(\Omega)$.

The pair (a_0, b_0) defines a self-adjoint linear operator $A_0: D(A_0) \subset L_s^2(\Omega) \rightarrow L_s^2(\Omega)$ with domain dense in $L_s^2(\Omega)$.

Furthermore, there are a nondecreasing sequence of eigenvalues $(\lambda_{0,j})_{j \in \mathbb{N}}$ and a corresponding L_s^2 -complete and L^2 -orthonormal system $(w_{0,j})_{j \in \mathbb{N}}$ of eigenvectors of the pair (a_0, b) (equivalently, of A_0).

It was proved in [15, Th. 3.3] that $\lambda_{\epsilon,j} \rightarrow \lambda_{0,j}$ as $\epsilon \rightarrow 0$ for $j \in \mathbb{N}$. Moreover, if $(\epsilon_n)_{n \in \mathbb{N}}$ is an arbitrary sequence of positive numbers converging to 0, then the orthonormal system $(w_{0,j})_{j \in \mathbb{N}}$ can be chosen in such a way that, up to a subsequence, $|w_{\epsilon_n,j} - w_{0,j}|_{\epsilon_n} \rightarrow 0$ as $n \rightarrow \infty$ for $j \in \mathbb{N}$.

In the sequel we will work with a fixed domain and we will sometimes write L^p for $L^p(\Omega)$ and H^1 for $H^1(\Omega)$.

Now let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a given nonlinearity. We will discuss some conditions on f which guarantee that the Nemitski operator $\hat{f}$ defined, for $u: \Omega \rightarrow \mathbb{R}$, by $\hat{f}(u) = f \circ u$ restricts to a C^1 -operator between certain function spaces.

Let us first recall the following regularity result for Nemitski operators, which is a slight modification of [1, Thms 3.4 and 3.7].

Theorem 2.1. *Let $\Omega \subset \mathbb{R}^n$ be open and bounded. Given $\rho, \sigma \in \mathbb{R}$ with $1 \leq \rho < \sigma$ set $\beta := (\sigma/\rho) - 1$ (observe $\beta > 0$). Suppose $f \in C^1(\mathbb{R} \rightarrow \mathbb{R})$ satisfies*

$$(2.1) \quad |f'(s)| \leq C(1 + |s|^\beta) \quad \text{for some } C \in]0, \infty[\text{ and all } s \in \mathbb{R}.$$

Then the Nemitski operator

$$\hat{f}: L^\sigma(\Omega) \rightarrow L^\rho(\Omega), \quad u \mapsto f \circ u$$

is well-defined and of class C^1 ; the Fréchet differential of $\hat{f}$ is given by $D\hat{f}(u)v = (f' \circ u) \cdot v$ for all u and $v \in L^\sigma(\Omega)$. Finally, there exists a constant $\hat{C} \in [0, \infty[$ such that the following estimates hold:

$$\begin{aligned} |\hat{f}(u)|_{L^\rho} &\leq \hat{C}(1 + |u|_{L^\sigma}^{\beta+1}) \\ \|D\hat{f}(u)\|_{\mathcal{L}(L^\sigma, L^\rho)} &\leq \hat{C}(1 + |u|_{L^\sigma}^\beta) \end{aligned}$$

for all $u \in L^\sigma(\Omega)$.

Theorem 2.1 implies the following

Corollary 2.2. *Let Ω be an open, bounded subset of $\mathbb{R}^2$, with Lipschitz boundary. Let $f \in C^1(\mathbb{R} \rightarrow \mathbb{R})$ satisfy the growth estimate*

$$|f'(s)| \leq C(1 + |s|^\beta) \quad \text{for } s \in \mathbb{R}$$

where C and $\beta \in [0, \infty[$ are arbitrary real constants. Let $F(y) := \int_0^y f(s) ds$ for $y \in \mathbb{R}$. Then $f \circ u \in L^2(\Omega)$ whenever $u \in H^1(\Omega)$. Moreover, the Nemitski operator

$$\hat{f}: H^1(\Omega) \rightarrow L^2(\Omega), \quad u \mapsto f \circ u,$$

is well-defined and of class C^1 on $H^1(\Omega)$; the Fréchet differential of $\hat{f}$ is given by $D\hat{f}(u)[v] = (f' \circ u) \cdot v$ for all u and $v \in H^1(\Omega)$. Let $\hat{f}|_{H_s^1}$ be the restriction of

$\hat{f}$ to $H_s^1(\Omega)$. Then $\hat{f}|_{H_s^1}$ is a C^1 map from $H_s^1(\Omega)$ to $L_s^2(\Omega)$, with $D(\hat{f}|_{H_s^1})(u) = (D\hat{f}(u))|_{H_s^1}$ for $u \in H_s^1(\Omega)$. Furthermore, $F \circ u \in L^1(\Omega)$ whenever $u \in H^1(\Omega)$ and the Nemitski operator

$$\hat{F}: H^1(\Omega) \rightarrow L^1(\Omega), \quad u \mapsto F \circ u,$$

is well-defined and of class C^1 on $H^1(\Omega)$; the Fréchet differential of $\hat{F}$ is given by $D\hat{F}(u)[v] = (f \circ u) \cdot v$ for all u and $v \in H^1(\Omega)$. Finally, there exists a constant $\hat{C}$ such that the following estimates hold:

$$\begin{aligned} |\hat{f}(u)|_{L^2} &\leq \hat{C}(1 + |u|_{H^1}^{\beta+1}) \\ \|D\hat{f}(u)\|_{\mathcal{L}(H^1, L^2)} &\leq \hat{C}(1 + |u|_{H^1}^\beta) \\ |\hat{F}(u)|_{L^1} &\leq \hat{C}(1 + |u|_{H^1}^{\beta+2}) \\ \|D\hat{F}(u)\|_{\mathcal{L}(H^1, L^1)} &\leq \hat{C}(1 + |u|_{H^1}^{\beta+1}) \end{aligned}$$

Proof. Without loss of generality we can assume that $\beta > 0$. Set $\rho = 2$ and $\sigma = 2(\beta + 1)$ (resp. $\rho = 1$ and $\sigma = \beta + 2$) in Theorem 2.1. By Sobolev's inequality,

$$|w|_{L^\sigma} \leq C_\sigma |w|_{H^1} \quad \text{for } w \in H^1(\Omega),$$

where C_σ is a real positive constant. Thus an application of Theorem 2.1 shows that the Nemitski operator $\hat{f}$ (resp. $\hat{F}$) is well-defined, of class C^1 and that the above estimates hold. We already know that $\hat{f}(u) \in L_s^2(\Omega)$ whenever $u \in H_s^1(\Omega)$ (Theorem 5.3 in [15]). In order to complete the proof, we only need to show that, if u and $v \in H_s^1(\Omega)$, then $D\hat{f}(u)[v] \in L_s^2(\Omega)$. This follows immediately from the formula

$$D\hat{f}(u)[v] = \lim_{t \rightarrow 0} \frac{\hat{f}(u + tv) - \hat{f}(u)}{t}$$

and from the fact that $L_s^2(\Omega)$ is closed in $L^2(\Omega)$. $\square$

For every $\epsilon > 0$ the linear operator A_ϵ is sectorial on $X = L^2(\Omega)$ and the fractional power space $X^{1/2}$ is equal to $H^1(\Omega)$. Similarly, A_0 is sectorial on $X = L_s^2(\Omega)$ and in this case $X^{1/2} = H_s^1(\Omega)$.

Let $g: H^1(\Omega) \rightarrow L^2(\Omega)$ be Lipschitzian on the bounded subsets of $H^1(\Omega)$ and such that $g(H_s^1(\Omega)) \subset L_s^2(\Omega)$. It follows from the results of [10] that for every $\epsilon > 0$ there exists a well-defined local semiflow $\pi_{\epsilon, g}$ on $H^1(\Omega)$ generated by the semilinear parabolic equation

$$(2.2) \quad \dot{u} + A_\epsilon u = g(u).$$

Moreover, there exists a well-defined local semiflow $\pi_{0, g}$ on $H_s^1(\Omega)$ generated by the semilinear parabolic equation

$$(2.3) \quad \dot{u} + A_0 u = g(u).$$

In particular, if a function $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies the assumptions of Corollary 2.2 then, by this corollary, $g := \hat{f}: H^1(\Omega) \rightarrow L^2(\Omega)$ is Lipschitzian on the bounded subsets of $H^1(\Omega)$ and $g(H_s^1(\Omega)) \subset L_s^2(\Omega)$, so the local semiflows $\pi_{\epsilon, \hat{f}}$, $\epsilon \geq 0$, are well-defined.

3. DESCRIPTION OF THE LIMIT PROBLEM

In our previous paper [15] we introduced the class of the so-called nicely decomposed domains. We showed that on nicely decomposed domains both the operator $A = A_0$ and its domain of definition can be explicitly characterized.

In this section we continue the study of nicely decomposed domains Ω and show, in particular, how to characterize the spaces $H_s^1(\Omega)$ and $L_s^2(\Omega)$ (Proposition 3.6). We also obtain a compact imbedding result (Proposition 3.4) which is important in establishing the gap condition (4.1) in section 3.

For the reader's convenience we will first recall the definition of a nicely decomposed domains.

We say that an open set $\Omega \subset \mathbb{R}^2$ *has connected vertical sections* if for every $x \in \mathbb{R}$ the x -section Ω_x is connected. Of course, this section is nonempty if and only if $x \in P(\Omega)$, where $P: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $(x, y) \mapsto x$ is the projection onto the first component. Note that, given a nonempty bounded domain Ω in $\mathbb{R}^2$, $J_\Omega := P(\Omega)$ is an open interval in $\mathbb{R}$, that is $J_\Omega =]a_\Omega, b_\Omega[$, where $-\infty < a_\Omega < b_\Omega < \infty$.

Given $a \in \mathbb{R}$ and $\delta \in]0, \infty[$ we set

$$I_\delta(a) :=]a - \delta, a + \delta[, \quad I_\delta^-(a) :=]a - \delta, a[\text{ and } I_\delta^+(a) :=]a, a + \delta[.$$

Definition 3.1. Let $\Omega, \Omega_1, \Omega_2$ be nonempty bounded domains in $\mathbb{R}^2$. Set $a_i := a_{\Omega_i}$ and $b_i := b_{\Omega_i}$, $i = 1, 2$. Given $c \in \mathbb{R}$ we say that Ω_1 *joins* Ω_2 *at* c *in* Ω if the following properties hold:

- (1) $\overline{\Omega}_1 \cap \overline{\Omega}_2 = \{c\} \times [\beta, \gamma]$ where $\beta = \beta_{\Omega_1, \Omega_2}$ and $\gamma = \gamma_{\Omega_1, \Omega_2}$ are some real numbers with $\beta < \gamma$;
- (2) $c = a_{\Omega_2} = b_{\Omega_1}$;
- (3) $\{c\} \times]\beta, \gamma[\subset \Omega$;
- (4) whenever $d \in]\beta, \gamma[$, then there is a $\delta = \delta(d) > 0$ with the property that

$$I_\delta(d) \subset]\beta, \gamma[$$

and

$$I_\delta^-(c) \times I_\delta(d) \subset \Omega_1, \quad I_\delta^+(c) \times I_\delta(d) \subset \Omega_2.$$

We say that Ω_1 and Ω_2 *join at* c *in* Ω if Ω_1 joins Ω_2 at c in Ω or Ω_2 joins Ω_1 at c in Ω .




Figure 1

Definition 3.2, cf Figure 1. Assume that $\Omega \subset \mathbb{R} \times \mathbb{R}$ is a nonempty bounded open domain with Lipschitz boundary. Let $P: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $(x, y) \mapsto x$ be the projection onto the first variable. A *nice decomposition* of Ω is a collection $\Omega_1, \dots, \Omega_r$ of nonempty pairwise disjoint open connected subsets of Ω with connected vertical sections such that, defining $J_k := J_{\Omega_k}$, $a_k := a_{\Omega_k}$, $b_k := b_{\Omega_k}$, $k = 1, \dots, r$, the following properties are satisfied:

- (1) $\Omega \setminus (\bigcup_{k=1}^r \Omega_k) \subset Z$, where $Z := \bigcup_{l=1}^r (\{a_l, b_l\} \times \mathbb{R})$;
- (2) whenever $k = 1, \dots, r$ then $\partial\Omega_k \subset \partial\Omega \cup (\{a_k, b_k\} \times \mathbb{R})$ and for $c \in \{a_k, b_k\}$ $\partial\Omega_k \cap (\{c\} \times \mathbb{R}) = \{c\} \times I$, where I is a compact (possibly degenerate) interval in $\mathbb{R}$;
- (3) whenever $k, l = 1, \dots, r$, $k \neq l$ and $(c, d) \in \overline{\Omega_k} \cap \overline{\Omega_l}$ is arbitrary then either Ω_k and Ω_l join at c in Ω or else there is an $m \in \{1, \dots, r\}$ such that Ω_k and Ω_m join at c in Ω and Ω_l and Ω_m join at c in Ω ;
- (4) for every $k = 1, \dots, r$ the function $p_k: J_k \rightarrow]0, \infty[$, $x \mapsto \mathcal{L}^1((\Omega_k)_x)$, is such that $1/p_k \in L^1(J_k)$.

Remarks.

- (1) Definition 3.2 says that, up to a set of measure zero, contained in a set Z of finitely many vertical lines, Ω can be decomposed into the finitely many domains Ω_k , $k = 1, \dots, r$ in such a way that at Z the various sets Ω_k and

- Ω_l ‘join’ in a nice way. Points of $\overline{\Omega} \cap Z$ are, intuitively speaking, those at which connected components of the vertical sections Ω_x bifurcate.
- (2) Let $R_1, \dots, R_s$ be closed bounded rectangles in $\mathbb{R}^2$ with edges parallel to the coordinate axes, and Ω_R be the interior of the union $\bigcup_{k=1}^s R_k$. Then, clearly, every connected component of Ω_R is a nicely decomposable domain with Lipschitz boundary.
 - (3) Let $\phi: \mathbb{R}^2 \rightarrow \mathbb{R}$ be a C^2 function, such that $\phi(x, y) \rightarrow \infty$ as $|(x, y)| \rightarrow \infty$. Assume that 0 is a regular value of ϕ . Moreover, suppose that, whenever $\phi(x_0, y_0) = 0$ and $\phi_y(x_0, y_0) = 0$ then $\phi_{yy}(x_0, y_0) \neq 0$. Let $\Omega_\phi := \{(x, y) \mid \phi(x, y) < 0\}$ and let Ω be any connected component of Ω_ϕ . It is a nice exercise to check that Ω is nicely decomposable.
 - (4) It can be shown that all real analytic domains are nicely decomposable.
- Finally, given a nice decomposition $\Omega_1, \dots, \Omega_r$ of Ω , we set

$$E := \bigcup_{k=1}^r ((\{a_k, b_k\} \times \mathbb{R}) \cap \partial\Omega_k).$$

Our goal is to give a detailed description of the spaces $H_s^1(\Omega)$ and $L_s^2(\Omega)$ when Ω is a nicely decomposed domain. We begin our description by first considering the simpler case of an open set O with connected vertical sections. Below, such a role will be played by the sets Ω_k occurring in the nice decomposition of Ω . We do not assume that O has Lipschitz boundary, since the sets Ω_k occurring in the nice decomposition of Ω in general do not have this property.

Let $P: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $(x, y) \mapsto x$ be the projection onto the first variable. Let $J := P(O)$ and assume for simplicity that $J =]0, 1[$.

The following proposition is proved in [15]:

Proposition 3.3. *Suppose O has connected vertical sections. Let $J := P(O)$ and define the function $p: J \rightarrow]0, \infty[$ by $x \mapsto \mathcal{L}^1(\Omega_x)$. If $u \in L^2(O)$ satisfies $u_y = 0$ in the distributional sense, then there is a set S of measure 0 in $\mathbb{R}^2$ and a function $v \in L_{\text{loc}}^1(J)$ such that $u(x, y) = v(x)$ for every $(x, y) \in O \setminus S$. Moreover, $p^{1/2}v \in L^2(J)$. If $u \in H^1(O)$ then $v' \in L_{\text{loc}}^1(J)$ and we can choose the set S so that $u(x, y) = v(x)$ and $u_x(x, y) = v'(x)$ for every $(x, y) \in \Omega \setminus S$. Moreover, $p^{1/2}v' \in L^2(J)$ and we can choose the function v to be absolutely continuous on J . The function $\tilde{u}: O \rightarrow \mathbb{R}$, $\tilde{u}(x, y) = v(x)$ is then a continuous representative of u . $\square$*

Now, since O is open and bounded, it is easy to check that the function p satisfies the following property:

(A) $p \in L^\infty(0, 1)$ and for every ϵ , $0 < \epsilon < 1 - \epsilon$, there exists $\delta > 0$ such that $p(x) > \delta$ a.e. in $]\epsilon, 1 - \epsilon[$.

Now given an arbitrary function p satisfying property (A) note that $p(x) > 0$ a.e. in $]0, 1[$. Therefore the linear spaces

$$H(p) := \left\{ u \in L_{\text{loc}}^1(0, 1) \mid p^{1/2}u \in L^2(0, 1) \right\}$$

and

$$V(p) := \left\{ u \in L^1_{\text{loc}}(0, 1) \mid u' \in L^1_{\text{loc}}(0, 1), p^{1/2}u \in L^2(0, 1), p^{1/2}u' \in L^2(0, 1) \right\}$$

are well-defined. Define on $H(p)$ and $V(p)$ the scalar products

$$\langle u, v \rangle_{H(p)} := \int_0^1 p(x)u(x)v(x) \, dx$$

and

$$\langle u, v \rangle_{V(p)} := \int_0^1 p(x)u'(x)v'(x) \, dx + \int_0^1 p(x)u(x)v(x) \, dx.$$

It is easy to check that these products define Hilbert space structures on $H(p)$ and $V(p)$.

Now, given O and p as in Proposition 3.3, define the mapping

$$\iota: L^2_s(O) \rightarrow H(p), \quad u \mapsto v,$$

where v is the function given by Proposition 3.3. It turns out that ι is an isometry of $L^2_s(O)$ onto $H(p)$. Moreover, ι restricts to an isometry of $H^1_s(O)$ onto $V(p)$.

We have the following result:

Proposition 3.4. *Assume that the function p satisfies (A) and $(1/p) \in L^1(0, 1)$. Let $u \in V(p)$. Then there exists a function $v \in C^0([0, 1])$ such that $u = v$ a.e. in $]0, 1[$. Moreover, the imbedding $V(p) \hookrightarrow C^0([0, 1])$ (and hence the imbedding $V(p) \hookrightarrow H(p)$) is compact.*

Proof. Let ϵ , $0 < \epsilon < 1 - \epsilon$, be arbitrary. Then there exists $\delta > 0$ such that $p(x) \geq \delta$ a.e. in $]\epsilon, 1 - \epsilon[$. It follows that $u|_{] \epsilon, 1 - \epsilon[} \in H^1(\epsilon, 1 - \epsilon)$. Since ϵ is arbitrary, we obtain that there exists a function $v \in C^0([0, 1])$ such that $u = v$ a.e. in $]0, 1[$. We show that v can be extended to a continuous function on $[0, 1]$. In fact, if x and $x' \in]0, 1[$, $x < x'$, then, by Hölder's inequality,

$$|v(x') - v(x)|^2 = \left| \int_x^{x'} p^{-1/2} p^{1/2} v' \, ds \right|^2 \leq \int_x^{x'} (1/p) \, ds \int_x^{x'} p v'^2 \, ds.$$

It follows that

$$(3.1) \quad |v(x') - v(x)| \leq |v|_{V(p)} \left(\int_x^{x'} (1/p) \, ds \right)^{1/2}.$$

Since $(1/p) \in L^1(0, 1)$, then $|v(x') - v(x)| \rightarrow 0$ as $x, x' \rightarrow 0$ (resp. $x, x' \rightarrow 1$), so there exist the limits $v(0) := \lim_{x \rightarrow 0} v(x)$ and $v(1) := \lim_{x \rightarrow 1} v(x)$. By continuity, estimate (3.1) holds for x and $x' \in [0, 1]$. Now fix $\bar{\epsilon}$, $0 < \bar{\epsilon} < 1 - \bar{\epsilon}$. Since $v|_{] \bar{\epsilon}, 1 - \bar{\epsilon}[} \in H^1(\bar{\epsilon}, 1 - \bar{\epsilon})$, it follows that there exists a constant K_1 such that

$$|v|_{L^\infty(\bar{\epsilon}, 1 - \bar{\epsilon})} \leq K_1 |v|_{H^1(\bar{\epsilon}, 1 - \bar{\epsilon})}.$$

Then we can find another constant K_2 , such that

$$(3.2) \quad |v|_{L^\infty(\bar{\epsilon}, 1-\bar{\epsilon})} \leq K_2 |v|_{V(p)}.$$

Now (3.1) and (3.2) together, imply that there is a third constant K_3 such that

$$(3.3) \quad |v|_{L^\infty(0,1)} \leq K_3 |v|_{V(p)},$$

so $V(p) \hookrightarrow C^0([0,1])$ with continuous imbedding. Finally, (3.3) and (3.1), and a straightforward application of Ascoli-Arzelà theorem, imply that the imbedding is compact. $\square$

Now we consider the full nicely decomposed domain Ω .

Lemma 3.5. *Let Ω be a nicely decomposed domain. Then, for $k = 1, \dots, r$, the following properties hold:*

- (1) *whenever $u \in L_s^2(\Omega)$, then $u|_{\Omega_k} \in L_s^2(\Omega_k)$;*
- (2) *whenever $u \in H_s^1(\Omega)$, then $u|_{\Omega_k} \in H_s^1(\Omega_k)$.*

Proof. Part (2) is obvious and part (1) follows directly from part (2) and from the definition of $L_s^2(\Omega)$ and $L_s^2(\Omega_k)$. $\square$

For $k = 1, \dots, r$ let us define the linear spaces

$$H_k := \left\{ u \in L_{\text{loc}}^1(a_k, b_k) \mid p_k^{1/2} u \in L^2(a_k, b_k) \right\}$$

and

$$V_k := \left\{ u \in H_k \mid u' \in L_{\text{loc}}^1(a_k, b_k), p_k^{1/2} u' \in L^2(a_k, b_k) \right\}.$$

We have seen that the spaces H_k and V_k endowed with the scalar products

$$\langle u, v \rangle_{H_k} := \int_{a_k}^{b_k} p_k(x) u(x) v(x) \, dx$$

and

$$\langle u, v \rangle_{V_k} := \int_{a_k}^{b_k} p_k(x) u'(x) v'(x) \, dx + \int_{a_k}^{b_k} p_k(x) u(x) v(x) \, dx$$

respectively, are Hilbert spaces and that the imbedding $V_k \hookrightarrow H_k$ is dense and compact. Moreover, consider the following bilinear forms on V_k :

$$a_{\langle k \rangle}(u, v) := \int_{a_k}^{b_k} p_k(x) u'(x) v'(x) \, dx,$$

$$b_{\langle k \rangle}(u, v) := \int_{a_k}^{b_k} p_k(x) u(x) v(x) \, dx.$$

Define the product spaces

$$H_{\oplus} := H_1 \oplus \cdots \oplus H_r := \{ [u] = (u_1, \dots, u_r) \mid u_k \in H_k, k = 1, \dots, r \}$$

and

$$V_{\oplus} := V_1 \oplus \cdots \oplus V_r := \{ [u] = (u_1, \dots, u_r) \mid u_k \in V_k, k = 1, \dots, r \},$$

with the scalar products

$$\langle [u], [v] \rangle_{H_{\oplus}} := \sum_{k=1}^r \langle u_k, v_k \rangle_{H_k}$$

and

$$\langle [u], [v] \rangle_{V_{\oplus}} := \sum_{k=1}^r \langle u_k, v_k \rangle_{V_k}$$

respectively. It is easy to check that $H_{\oplus}$ and $V_{\oplus}$ are Hilbert spaces and that the imbedding $V_{\oplus} \hookrightarrow H_{\oplus}$ is dense and compact.

Furthermore, consider the following bilinear forms on $V_{\oplus}$:

$$a_{\oplus}([u], [v]) := \sum_{k=1}^r a_{\langle k \rangle}(u_k, v_k)$$

and

$$b_{\oplus}([u], [v]) := \sum_{k=1}^r b_{\langle k \rangle}(u_k, v_k).$$

Note that $b_{\oplus}$ is just the restriction to $V_{\oplus} \times V_{\oplus}$ of the scalar product $\langle \cdot, \cdot \rangle_{H_{\oplus}}$.

For $k = 1, \dots, r$, let us define the mapping

$$\iota_k: L_s^2(\Omega_k) \rightarrow H_k, \quad u \mapsto v,$$

where v is the function given by Proposition 3.3. It turns out that ι_k is an isometry of $L_s^2(\Omega_k)$ onto H_k and that ι_k restricts to an isometry of $H_s^1(\Omega_k)$ onto V_k . Moreover, let us define

$$\iota_{\oplus}: L_s^2(\Omega) \rightarrow H_{\oplus}, \quad \iota_{\oplus} u := (\iota_1(u|_{\Omega_1}), \dots, \iota_r(u|_{\Omega_r})).$$

It follows that $\iota_{\oplus}$ is an isometry of $L_s^2(\Omega)$ into $H_{\oplus}$ and that $\iota_{\oplus}$ restricts to an isometry of $H_s^1(\Omega)$ into $V_{\oplus}$.

Define

$$V_{\oplus}^{\sim} := \left\{ [u] \in V_{\oplus} \mid u_k(b_k) = u_l(a_l) \text{ whenever } b_k = a_l = c \text{ and } \Omega_k \text{ and } \Omega_l \text{ join at } c. \right\}$$

Now we are able to characterize the spaces $H_s^1(\Omega)$ and $L_s^2(\Omega)$:

Proposition 3.6. *The following properties hold:*

- (1) $\iota_{\oplus}(L_s^2(\Omega)) = H_{\oplus};$
- (2) $\iota_{\oplus}(H_s^1(\Omega)) = V_{\oplus}^{\sim}.$

Proof. This result is more or less implicitly contained in the proofs of Theorems 6.5 and 6.6 in [15]. For the reader's convenience, we summarize the main steps of the proof.

First we prove (2). Let $u \in H_s^1(\Omega)$. We shall prove that $\iota_{\oplus}u \in V_{\oplus}^{\sim}$. Assume that for some $k, l \in \{1, \dots, r\}$, $b_k = a_l = c$ and Ω_k and Ω_l join at c . We have to prove that

$$\iota_k(u|_{\Omega_k})(c) = \iota_l(u|_{\Omega_l})(c).$$

Set

$$v_k := \iota_k(u|_{\Omega_k}), \quad v_l := \iota_l(u|_{\Omega_l}).$$

Then, by Lemma 6.5 in [15], we immediately obtain that $v_k(c) = v_l(c)$, and we are done.

Now assume, conversely, that $[u] \in V_{\oplus}^{\sim}$. We shall prove that there exists $v \in H_s^1(\Omega)$ such that $\iota_{\oplus}v = [u]$. We define a function v on Ω in the following way:

$$v(x, y) := \begin{cases} u_k(x) & \text{for all } (x, y) \in \Omega_k, k = 1, \dots, r \\ 0 & \text{for all } (x, y) \in \Omega \setminus (\bigcup_{k=1}^r \Omega_k). \end{cases}$$

As in the proof of Theorem 6.6 in [15], one can show that $v \in H_s^1(\Omega)$ and that

$$\begin{aligned} v_x(x, y) &= u'_k(x) && \text{a.e. in } \Omega_k, k = 1, \dots, r, \\ v_y(x, y) &= 0 && \text{a.e. in } \Omega. \end{aligned}$$

Finally observe that, by construction, $\iota_{\oplus}v = [u]$ and the proof of (2) is complete.

Now we prove (1). Let $[u] \in H_{\oplus}$. We shall prove that there exists $v \in L_s^2(\Omega)$ such that $[u] = \iota_{\oplus}v$. We define a function v on Ω in the following way:

$$v(x, y) := \begin{cases} u_k(x) & \text{for all } (x, y) \in \Omega_k, k = 1, \dots, r \\ 0 & \text{for all } (x, y) \in \Omega \setminus (\bigcup_{k=1}^r \Omega_k) \end{cases}$$

Then $v \in L^2(\Omega)$. We claim that $v \in L_s^2(\Omega)$. This means that v can be approximated in the $L^2(\Omega)$ norm by functions of $H_s^1(\Omega)$. Let $\epsilon > 0$ be arbitrary. For $k = 1, \dots, r$, take $\phi_k \in C_0^\infty([a_k, b_k])$ such that

$$\int_{a_k}^{b_k} p_k |u_k - \phi_k|^2 dx < \frac{\epsilon}{k}.$$

Define a function ψ on Ω in the following way:

$$\psi(x, y) := \begin{cases} \phi_k(x) & \text{for all } (x, y) \in \Omega_k, k = 1, \dots, r \\ 0 & \text{for all } (x, y) \in \Omega \setminus (\bigcup_{k=1}^r \Omega_k) \end{cases}$$

By the proof of (2), $\psi \in H_s^1(\Omega)$. Moreover,

$$\int_{\Omega} |v - \psi|^2 dx dy = \sum_{k=1}^r \int_{\Omega_k} |v - \psi|^2 dx dy = \sum_{k=1}^r \int_{a_k}^{b_k} p_k |u_k - \phi_k|^2 dx < \epsilon,$$

and our claim is proved. Finally observe that, by definition, $\iota_{\oplus} v = [u]$ and the proof is complete. $\square$

We end this section with some remarks concerning the self-adjoint operator A generated by a in $L_s^2(\Omega)$. Let $a_{\oplus}$ be the restriction of $a_{\oplus}$ to $V_{\oplus} \times V_{\oplus}$ and let $A_{\oplus}$ be the self-adjoint operator generated by $a_{\oplus}$ in $H_{\oplus}$. If $u \in D(A)$, then, for all $v \in H_s^1(\Omega)$,

$$\langle Au, v \rangle_{L_s^2(\Omega)} = a(u, v) = a_{\oplus}(\iota_{\oplus} u, \iota_{\oplus} v).$$

On the other hand,

$$\langle Au, v \rangle_{L_s^2(\Omega)} = \langle \iota_{\oplus} Au, \iota_{\oplus} v \rangle_{H_{\oplus}}.$$

It follows that

$$a_{\oplus}(\iota_{\oplus} u, \iota_{\oplus} v) = \langle \iota_{\oplus} Au, \iota_{\oplus} v \rangle_{H_{\oplus}}$$

for all $v \in H_s^1(\Omega)$, so $\iota_{\oplus} u \in D(A_{\oplus})$ and $A_{\oplus} \iota_{\oplus} u = \iota_{\oplus} Au$. Similarly, one can prove that, whenever $[u] \in D(A_{\oplus})$, then $\iota_{\oplus}^{-1}[u] \in D(A)$, and $A \iota_{\oplus}^{-1}[u] = \iota_{\oplus}^{-1} A_{\oplus}[u]$. This means that $\iota_{\oplus}$ restricts to an isometry of $D(A)$ onto $D(A_{\oplus})$ and that $A = \iota_{\oplus}^{-1} A_{\oplus} \iota_{\oplus}$.

For $k = 1, \dots, r$, let us define the spaces

$$Z_k := \left\{ u \in V_k \mid (p_k u')' \in L_{\text{loc}}^1(a_k, b_k), p_k^{-1/2} (p_k u')' \in L^2(a_k, b_k) \right\},$$

and let

$$Z_{\oplus} := Z_1 \oplus \dots \oplus Z_r.$$

It is not difficult to show that, whenever $u \in Z_k$, then $p_k u'$ can be extended to a continuous function on $[a_k, b_k]$ (cf. the proof of Proposition 3.4). Now we can restate Theorem 6.6 in [15] in the following way:

Theorem 3.7. *Let $A_{\oplus}$ be the self-adjoint operator generated by the bilinear form $a_{\oplus}$. Then $D(A_{\oplus}) = Z_{\oplus}$, where $Z_{\oplus}$ is the subspace of $Z_{\oplus}$ consisting of all $[u] = (u_1, \dots, u_k)$ satisfying the following properties:*

- (1) $u_k(b_k) = u_l(a_l)$ whenever $b_k = a_l = c$ and Ω_k and Ω_l join at c ;
- (2) whenever Γ is a connected component of E (necessarily of the form $\Gamma = \{c\} \times I$, where $c \in \bigcup_{k=1}^r \{a_k, b_k\}$ and I is an interval) then

$$\sum_{k \in \sigma_+} (p_k u_k')(c) = \sum_{k \in \sigma_-} (p_k u_k')(c).$$

Here, $\sigma_+ = \sigma_+(\Gamma)$ is the set of all k such that $\overline{\Omega}_k \cap (\{b_k\} \times \mathbb{R}) \subset \Gamma$ and so $b_k = c$, while $\sigma_- = \sigma_-(\Gamma)$ is the set of all k such that $\overline{\Omega}_k \cap (\{a_k\} \times \mathbb{R}) \subset \Gamma$ and so $a_k = c$.

Moreover, for $[u] \in Z_{\oplus}^{\sim}$, $A_{\oplus}^{\sim}[u] = (p_1^{-1}(p_1 u'_r)', \dots, p_r^{-1}(p_r u'_r)'). \quad \square$

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a C^1 -function, satisfying the assumptions of Corollary 2.2. As in Section 2 consider the abstract semilinear parabolic equation

$$(3.4) \quad \dot{u} = Au + \hat{f}(u)$$

on $H_s^1(\Omega)$. Assume that Ω is nicely decomposable. Then, due to the isometry $\iota_{\oplus}$ and in view of Theorem 3.7, the abstract equation (3.4) is equivalent to the following system of ‘concrete’ one-dimensional reaction-diffusion equations:

$$\partial_t u_k = (1/p_k)(p_k u_k')' + f(u_k) \quad \text{on }]a_k, b_k[, \text{ for } k = 1, \dots, r,$$

with compatibility conditions

$$u_k(c) = u_l(c)$$

whenever $b_k = a_l = c$ and Ω_k and Ω_l join at c , and Kirchhoff type balance conditions

$$\sum_{k \in \sigma_+(\Gamma)} (p_k u_k')(c) = \sum_{k \in \sigma_-(\Gamma)} (p_k u_k')(c)$$

whenever $\Gamma = \{c\} \times I$ is a connected component of E .




Figure 2

As it was explained in [15], such a system can be interpreted as a reaction-diffusion equation on an appropriate finite graph $\mathcal{G}$. Figure 1 suggests that the edges of this graph are the intervals $[a_k, b_k]$, for $k = 1, \dots, r$ and their endpoints are its vertices. Moreover, each interval should be repeated the number of times it occurs in the sequence $([a_k, b_k])_{k=1}^r$. On the other hand each endpoint c should be repeated the number of times it occurs as the x -component of a connected component of E (see Figure 2).

4. SPECTRAL GAP CONDITION

In this section let $\Omega \subset \mathbb{R}^2$ be a nonempty bounded open set with Lipschitz boundary. We denote by $(\lambda_\nu)_{\nu \in \mathbb{N}}$ the repeated sequence of the eigenvalues of the pair (a, b) where

$$a(u, v) := \int_{\Omega} \nabla u \cdot \nabla v \, dx \, dy = \int_{\Omega} u_x v_x \, dx \, dy \quad u, v \in H_s^1(\Omega)$$

and

$$b(u, v) = \int_{\Omega} uv \, dx \, dy \quad u, v \in H_s^1(\Omega).$$

We will prove in this section that, under some additional assumptions on Ω (cf. condition (C) below), the following so-called *gap condition* holds:

$$(4.1.) \quad \limsup_{\nu \rightarrow \infty} \frac{\lambda_{\nu+1} - \lambda_{\nu}}{\lambda_{\nu}^{1/2}} > 0,$$

This gap condition will enable us, in the next section, to establish the existence of inertial manifolds for the semiflows π_{ϵ} , for $\epsilon \geq 0$ small enough.

We first prove a simple general estimate on the sequence (λ_{ν}) :

Proposition 4.1. *There is a $\beta \in]0, \infty[$ such that*

$$\lambda_{\nu} \leq \beta^2 \nu^2 \quad \text{for all } \nu \in \mathbb{N}.$$

Proof. There are nonempty and bounded open intervals I, I', J and $J' \subset \mathbb{R}$ such that

$$I \times J \subset \Omega \subset I' \times J'.$$

Let γ, γ' and δ be the lengths of J, J' and I , respectively. For $\nu \in \mathbb{N}$ let $\mathcal{V}_{\nu}$ (resp. $\tilde{\mathcal{V}}_{\nu}$) be the set of all ν -dimensional subspaces of $H_s^1(\Omega)$ (resp. $H_0^1(I)$). For every function $v \in H_0^1(I)$ consider its trivial extension $\tilde{v}: I' \rightarrow \mathbb{R}$, $\tilde{v}(x) := v(x)$ for $x \in I$ and $\tilde{v}(x) := 0$ for $x \in I' \setminus I$. It follows that $\tilde{v} \in H_0^1(I')$. Therefore the function $u: I' \times J' \rightarrow \mathbb{R}$ defined by $u(x, y) = \tilde{v}(x)$ for $(x, y) \in I' \times J'$ lies in $H^1(I' \times J')$, so $u|_{\Omega} \in H^1(\Omega)$. By Theorem 2.5 in [15] we have $u \in H_s^1(\Omega)$. The operator

$$\Phi: H_0^1(I) \rightarrow H_s^1(\Omega), \quad v \mapsto u,$$

is linear and injective. In particular, $\Phi(\tilde{\mathcal{V}}_{\nu}) \subset \mathcal{V}_{\nu}$, $\nu \in \mathbb{N}$. Thus, for every $\nu \in \mathbb{N}$,

$$\begin{aligned} \lambda_{\nu} &= \inf_{E \in \mathcal{V}_{\nu}} \sup_{u \in E \setminus \{0\}} \frac{a(u, u)}{b(u, u)} \leq \inf_{E \in \Phi(\tilde{\mathcal{V}}_{\nu})} \sup_{u \in E \setminus \{0\}} \frac{a(u, u)}{b(u, u)} \\ &= \inf_{E \in \tilde{\mathcal{V}}_{\nu}} \sup_{v \in E \setminus \{0\}} \frac{a(\Phi v, \Phi v)}{b(\Phi v, \Phi v)} \leq \inf_{E \in \tilde{\mathcal{V}}_{\nu}} \sup_{v \in E \setminus \{0\}} \frac{\gamma' \int_I v' \cdot v' \, dx}{\gamma \int_I v \cdot v \, dx} = (\gamma'/\gamma) \mu_{\nu}, \end{aligned}$$

where μ_{ν} are the eigenvalues of the Dirichlet problem

$$\begin{cases} v''(x) = -\mu v(x), & x \in I \\ v(x) = 0, & x \in \partial I. \end{cases}$$

It follows that $\mu_{\nu} = (\pi/\delta)^2 \nu^2$, $\nu \in \mathbb{N}$. The proposition is proved. $\square$

For the rest of this section assume that Ω admits a nice decomposition $\Omega_1, \dots, \Omega_r$. Using the notation of the last section we obtain from Proposition 2.2 in [15] an eigenvalue-eigenvector sequence $(\lambda_\nu^k, u_\nu^k)_{\nu \in \mathbb{N}}$ of the pair (a_k, b_k) of bilinear forms such that

$$0 = \lambda_1^k \leq \lambda_2^k \leq \lambda_3^k \leq \dots$$

and $(u_\nu^k)_{\nu \in \mathbb{N}}$ is a complete orthonormal system in H_k , $k = 1, \dots, r$. We also obtain an eigenvalue-eigenvector sequence $(\lambda_\nu^\oplus, [u]_\nu^\oplus)_{\nu \in \mathbb{N}}$ of the pair $(a_\oplus, b_\oplus)$ such that

$$0 \leq \lambda_1^\oplus \leq \lambda_2^\oplus \leq \lambda_3^\oplus \leq \dots$$

and $([u]_\nu^\oplus)_{\nu \in \mathbb{N}}$ is a complete orthonormal system in $H_\oplus$.

The following result holds:

Proposition 4.2.

$$(4.2) \quad \lambda_\nu \geq \lambda_\nu^\oplus \quad \text{for all } \nu \in \mathbb{N}.$$

Proof. The proof is very similar to that of Proposition 4.1. For $\nu \in \mathbb{N}$ let $\mathcal{V}_\nu$ (resp. $\tilde{\mathcal{V}}_\nu$) be the set of all ν -dimensional subspaces of $V_\oplus$ (resp. $H_s^1(\Omega)$). Since the map $\Phi := \iota_\oplus|_{H_s^1(\Omega)}: H_s^1(\Omega) \rightarrow V_\oplus$ is an isometry (hence injective) it follows that $\Phi(\tilde{\mathcal{V}}_\nu) \subset \mathcal{V}_\nu$, $\nu \in \mathbb{N}$. Moreover, $a_\oplus(\Phi u, \Phi v) = a(u, v)$ and $b_\oplus(\Phi u, \Phi v) = b(u, v)$ for $u, v \in H_s^1(\Omega)$. Thus, for every $\nu \in \mathbb{N}$,

$$\begin{aligned} \lambda_\nu^\oplus &= \inf_{E \in \tilde{\mathcal{V}}_\nu} \sup_{[u] \in E \setminus \{0\}} \frac{a_\oplus([u], [u])}{b_\oplus([u], [u])} \leq \inf_{E \in \Phi(\tilde{\mathcal{V}}_\nu)} \sup_{[u] \in E \setminus \{0\}} \frac{a_\oplus([u], [u])}{b_\oplus([u], [u])} \\ &= \inf_{\tilde{E} \in \tilde{\mathcal{V}}_\nu} \sup_{u \in \tilde{E} \setminus \{0\}} \frac{a_\oplus(\Phi u, \Phi u)}{b_\oplus(\Phi u, \Phi u)} = \inf_{\tilde{E} \in \tilde{\mathcal{V}}_\nu} \sup_{u \in \tilde{E} \setminus \{0\}} \frac{a(u, u)}{b(u, u)} = \lambda_\nu. \end{aligned}$$

This completes the proof. $\square$

Now let p be a function satisfying the assumptions of Proposition 3.4. Consider the following bilinear form on $V(p)$:

$$a^p(u, v) := \int_0^1 p(x) u'(x) v'(x) \, dx.$$

Furthermore, define $b^p(u, v) := \langle u, v \rangle_{H(p)}$ for $u, v \in H(p)$. Then Proposition 3.4 and Proposition 2.2 in [15] implies that there exists a sequence $(\lambda_\nu^p, u_\nu^p)_{\nu \in \mathbb{N}}$ of eigenvalue-eigenvector pairs of (a^p, b^p) such that

$$0 = \lambda_1^p \leq \lambda_2^p \leq \lambda_3^p \leq \dots$$

and $(u_\nu^p)_{\nu \in \mathbb{N}}$ is a complete orthonormal system in $H(p)$.

The next proposition describes the asymptotic behavior of the sequence $(\lambda_\nu^p)_{\nu \in \mathbb{N}}$.

Proposition 4.3. *Let $p \in L^\infty(0, 1)$ satisfy one of the following conditions:*

- (1) *there exist two constants α and β , $0 < \alpha \leq \beta$, such that $\alpha \leq p(x) \leq \beta$ almost everywhere in $]0, 1[$;*
- (2) *there exists a function $q \in C^0([0, 1]) \cap C^2(]0, 1[)$, with $q(x) > 0$, $q'(x) \geq 0$ and $q''(x) \leq 0$ on $]0, 1[$, $q(0) = 0$ and $(1/q) \in L^1(0, 1)$, and there exist two constants α and β , $0 < \alpha \leq \beta$, such that $\alpha q(x) \leq p(x) \leq \beta q(x)$ almost everywhere in $]0, 1[$.*

Let $(\lambda_\nu^p)_{\nu \in \mathbb{N}}$ be the repeated sequence of the eigenvalues of (a^p, b^p) above. Then there exists a constant $\gamma > 0$, which depends only on α and β , such that

$$\lambda_\nu^p \geq \gamma(\nu - 1)^2, \quad \nu = 1, 2, \dots$$

The proof of Theorem 4.3 is based on the following

Lemma 4.4. *Let $q \in C^0([0, 1]) \cap C^2(]0, 1[)$, with $q(x) > 0$, $q'(x) \geq 0$ and $q''(x) \leq 0$ on $]0, 1[$, $q(0) = 0$ and $(1/q) \in L^1(0, 1)$. Let $(\lambda_\nu^q)_{\nu \in \mathbb{N}}$ be the repeated sequence of the eigenvalues of (a_q, b_q) above. Then*

$$\lambda_\nu^q \geq \pi^2(\nu - 1)^2, \quad \nu = 1, 2, \dots$$

Proof. See the Appendix. $\square$

Proof of Proposition 4.3. For every $\nu \in \mathbb{N}$,

$$\lambda_\nu^p = \inf_{E \in \mathcal{V}_\nu(p)} \sup_{u \in E \setminus \{0\}} \frac{a_p(u, u)}{b_p(u, u)},$$

where $\mathcal{V}_\nu(p)$ is the set of all ν -dimensional subspaces of $V(p)$. If condition (1) holds, then $H(p) = L^2(0, 1) = H(1)$ and $V(p) = H^1(0, 1) = V(1)$ with equivalent norms. Then

$$\begin{aligned} \lambda_\nu^p &= \inf_{E \in \mathcal{V}_\nu(p)} \sup_{u \in E \setminus \{0\}} \frac{\int_0^1 p u'^2 dx}{\int_0^1 p u^2 dx} \\ &\geq \inf_{E \in \mathcal{V}_\nu(1)} \sup_{u \in E \setminus \{0\}} \frac{\alpha \int_0^1 u'^2 dx}{\beta \int_0^1 u^2 dx} \\ &= (\alpha/\beta) \lambda_\nu^1, \quad \nu \in \mathbb{N} \end{aligned}$$

Now we observe that

$$\lambda_\nu^1 = \inf_{E \in \mathcal{V}_\nu(1)} \sup_{u \in E \setminus \{0\}} \frac{\int_0^1 u'^2 dx}{\int_0^1 u^2 dx}, \quad \nu = 1, 2, \dots$$

are the eigenvalues of the Neumann problem

$$\begin{cases} u''(x) = \lambda u(x), & x \in]0, 1[\\ u'(x) = 0, & x \in \{0, 1\} \end{cases}$$

This implies that $\lambda_\nu^1 = \pi^2(\nu-1)^2$, $\nu = 1, 2, \dots$, and the conclusion follows. Assume now that condition (2) holds. Then $H(p) = H(q)$ and $V(p) = V(q)$ with equivalent norms. Then

$$\begin{aligned} \lambda_\nu^p &= \inf_{E \in \mathcal{V}_\nu(p)} \sup_{u \in E \setminus \{0\}} \frac{\int_0^1 p u'^2 dx}{\int_0^1 p u^2 dx} \\ &\geq \inf_{E \in \mathcal{V}_\nu(q)} \sup_{u \in E \setminus \{0\}} \frac{\alpha \int_0^1 q u'^2 dx}{\beta \int_0^1 q u^2 dx} \\ &= (\alpha/\beta) \lambda_\nu^q, \quad \nu = 1, 2, \dots \end{aligned}$$

By Lemma 4.4, $\lambda_\nu^q \geq \pi^2(\nu-1)^2$, $\nu = 1, 2, \dots$, and the theorem is proved. $\square$

For $k = 1, \dots, r$ and $\tau \in \mathbb{R}$, $\tau > 0$, let us define

$$m(k, \tau) := \max \{ \nu \in \mathbb{N} \mid \lambda_\nu^k < \tau \},$$

and let

$$m(\oplus, \tau) := \max \{ \nu \in \mathbb{N} \mid \lambda_\nu^\oplus < \tau \}.$$

We have the following

Proposition 4.5. *Let $\tau \in \mathbb{R}$, $\tau > 0$. Then $m(\oplus, \tau) = \sum_{k=1}^r m(k, \tau)$.*

Proof. For $k = 1, \dots, r$ and $\nu \in \mathbb{N}$, define

$$[u]_\nu^k := (\underbrace{0, \dots, 0}_{k-1}, u_\nu^k, 0, \dots, 0).$$

It is easy to check that $(\lambda_\nu^k, [u]_\nu^k)$ is an eigenvalue-eigenvector pair of $(a_\oplus, b_\oplus)$ and that $([u]_\nu^k)_{\nu \in \mathbb{N}}^{k=1, \dots, r}$ is a complete orthonormal system in $H_\oplus$. The conclusion follows. $\square$

Assume now that Ω satisfies the following additional hypothesis:

(C) For every $k = 1, \dots, r$ one of the following conditions is satisfied:

- (1) there exist two constants α_k and β_k , $0 < \alpha_k \leq \beta_k$, such that $\alpha_k \leq p_k(x) \leq \beta_k$ in $]a_k, b_k[$;
- (2) there exists a function $q_k \in C^0([a_k, b_k]) \cap C^2(]a_k, b_k[)$, with $q_k(x) > 0$, $q'_k(x) \geq 0$ and $q''_k(x) \leq 0$ on $]a_k, b_k[$, $q_k(a_k) = 0$ and $(1/q_k) \in L^1(a_k, b_k)$, and there exist two constants α_k and β_k , $0 < \alpha_k \leq \beta_k$, such that $\alpha_k q_k(x) \leq p_k(x) \leq \beta_k q_k(x)$ in $]a_k, b_k[$;
- (3) there exists a function $q_k \in C^0([a_k, b_k]) \cap C^2([a_k, b_k])$, with $q_k(x) > 0$, $q'_k(x) \leq 0$ and $q''_k(x) \leq 0$ on $[a_k, b_k]$, $q_k(b_k) = 0$ and $(1/q_k) \in L^1(a_k, b_k)$, and there exist two constants α_k and β_k , $0 < \alpha_k \leq \beta_k$, such that $\alpha_k q_k(x) \leq p_k(x) \leq \beta_k q_k(x)$ in $]a_k, b_k[$.

Remark. The technical condition (C) is general enough to cover all the classes of nicely decomposable domains discussed in the remarks following Definition 3.2.

Proposition 4.6. *Assume Ω satisfies condition (C) above. Then*

$$\limsup_{\nu \rightarrow \infty} \frac{\lambda_{\nu}^{\oplus}}{\nu^2} > 0.$$

Proof. Condition (C), together with a simple renormalization of the interval $]a_k, b_k[$ and a straightforward application of Proposition 4.3, imply that for every $k = 1, \dots, r$ we can find a constant $\gamma_k > 0$ such that $\lambda_{\nu}^k \geq \gamma_k(\nu - 1)^2$ for $\nu = 1, 2, \dots$. Let $\gamma := \min\{\gamma_1, \dots, \gamma_r\}$. Then, for $k = 1, \dots, r$ and $\nu \in \mathbb{N}$, we have $m(k, \gamma(\nu - 1)^2) \leq \nu - 1$. It follows that $m(\oplus, \gamma(\nu - 1)^2) = \sum_{k=1}^r m(k, \gamma(\nu - 1)^2) \leq r(\nu - 1)$ for all $\nu \in \mathbb{N}$. But this means that $\lambda_{r(\nu-1)+1}^{\oplus} \geq \gamma(\nu - 1)^2$ for all $\nu \in \mathbb{N}$. Accordingly, we have:

$$\limsup_{\nu \rightarrow \infty} \frac{\lambda_{r(\nu-1)+1}^{\oplus}}{(r(\nu-1)+1)^2} \geq \limsup_{\nu \rightarrow \infty} \frac{\gamma(\nu-1)^2}{(r(\nu-1)+1)^2} = \frac{\gamma}{r^2} > 0$$

and the conclusion follows. $\square$

Proposition 4.7. *Let $(\mu_{\nu})_{\nu \in \mathbb{N}}$ be a nondecreasing sequence of nonnegative real numbers such that*

$$\limsup_{\nu \rightarrow \infty} \frac{\mu_{\nu}}{\nu^2} > 0$$

and

$$\mu_{\nu} \leq \beta^2 \nu^2 \quad \text{for some } \beta \in]0, \infty[\text{ and all } \nu \in \mathbb{N}.$$

Then

$$\limsup_{\nu \rightarrow \infty} \frac{\mu_{\nu+1} - \mu_{\nu}}{\mu_{\nu}^{1/2}} > 0.$$

Proof. Our hypotheses imply in particular, that $\mu_{\nu} > 0$ for all ν large enough. Thus, if the proposition is not true, then

$$\lim_{\nu \rightarrow \infty} \frac{\mu_{\nu+1} - \mu_{\nu}}{\mu_{\nu}^{1/2}} = 0.$$

Let $\epsilon > 0$ be arbitrary. Then there is a $\nu_0 \in \mathbb{N}$ such that

$$0 \leq \frac{\mu_{\nu+1} - \mu_{\nu}}{\mu_{\nu}^{1/2}} < \epsilon/\beta, \quad \nu \geq \nu_0.$$

Therefore

$$\mu_{\nu+1} \leq \mu_{\nu} + (\epsilon/\beta)\mu_{\nu}^{1/2} \leq \mu_{\nu} + \epsilon\nu, \quad \nu \geq \nu_0.$$

Hence,

$$(4.3) \quad \mu_{\nu+1} \leq \mu_{\nu_0} + \epsilon \left(\sum_{j=\nu_0}^{\nu} j \right) = \mu_{\nu_0} + (\epsilon/2) (\nu(\nu+1) - \nu_0(\nu_0+1)), \quad \nu \geq \nu_0.$$

Letting $\nu \rightarrow \infty$ in (4.3) implies that

$$\limsup_{\nu \rightarrow \infty} \frac{\mu_\nu}{\nu^2} \leq \frac{\epsilon}{2}.$$

Since $\epsilon > 0$ is arbitrary, it follows that

$$\limsup_{\nu \rightarrow \infty} \frac{\mu_\nu}{\nu^2} = 0,$$

a contradiction, which proves the proposition. $\square$

We can now state the main result of this section:

Theorem 4.8. *Let $\Omega \subset \mathbb{R}^2$ be a nicely decomposed domain and assume condition (C) is satisfied. Let $(\lambda_\nu)_{\nu \in \mathbb{N}}$ be the repeated sequence of eigenvalues of the pair (a, b) in $H_s^1(\Omega)$. Then the gap condition (4.1) is satisfied.*

Proof. Using Propositions 4.1 and 4.2 together with Proposition 4.6 we see that

$$\limsup_{\nu \rightarrow \infty} \frac{\lambda_\nu}{\nu^2} > 0$$

and

$$\lambda_\nu \leq \beta^2 \nu^2 \quad \text{for some } \beta \in]0, \infty[\text{ and all } \nu \in \mathbb{N}.$$

Therefore Proposition 4.7 completes the proof. $\square$

5. INERTIAL MANIFOLDS

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function satisfying the hypotheses of Corollary 2.2 together with the following dissipativeness condition:

$$\limsup_{|s| \rightarrow \infty} f(s)/s \leq -\delta_0 \quad \text{for some } \delta_0 > 0.$$

Then, by results of [15], for every $\epsilon \geq 0$ the semiflow $\pi_\epsilon := \pi_{\epsilon, \hat{f}}$ possesses a global attractor $\mathcal{A}_\epsilon$. Moreover, the family $(\mathcal{A}_\epsilon)_{\epsilon \geq 0}$ of attractors is upper semicontinuous at $\epsilon = 0$.

If the eigenvalues of the limit operator A_0 satisfy the gap condition (5.5) below, then, as we shall prove in Theorem 5.2 below, there exists an $\epsilon_0 > 0$ and a family $\mathcal{M}_\epsilon$, $0 \leq \epsilon \leq \epsilon_0$ of C^1 -manifolds of some finite dimension ν such that, whenever $0 \leq \epsilon \leq \epsilon_0$, then $\mathcal{A}_\epsilon \subset \mathcal{M}_\epsilon$ and the manifold $\mathcal{M}_\epsilon$ is locally invariant relative to the semiflow π_ϵ on a H^1 -neighborhood of the attractor $\mathcal{A}_\epsilon$. Moreover, the flows on the center manifolds $\mathcal{M}_\epsilon$ converge in the (regular) C^1 -sense to the flow on $\mathcal{M}_0$.

In particular, this result is valid on nicely decomposable domains satisfying condition (C) and so this extends and improves the corresponding results of Hale and Raugel [7] obtained for domains representable as ordinate sets with respect to some

positive function. In fact, as it was explained in the Introduction, the inertial manifolds constructed in [7] are relative to some modified semiflows, which are equal to the original semiflows π_ϵ on the attractor $\mathcal{A}_\epsilon$ but are different from π_ϵ on every neighborhood of it.

The proof of Theorem 5.2 is based on the method of functions of exponential growth, used before by many researchers (cf. [4], [18] and the references contained in these papers). First we choose an open set U in $H^1(\Omega)$ which includes all the attractors $\mathcal{A}_\epsilon$, $\epsilon \in [0, \epsilon_0]$, $\epsilon_0 > 0$ small. Then we modify the Nemitski operator $\hat{f}$ by finding a globally Lipschitzian map $g: H^1(\Omega) \rightarrow L^2(\Omega)$ with $\hat{f}(u) = g(u)$ for $u \in U$. For fixed $\epsilon \in [0, \epsilon_0]$, we seek an invariant manifold $\mathcal{M}_\epsilon$ for the modified semiflow $\pi_{\epsilon, g}$ in the form $\mathcal{M}_\epsilon = \Lambda_\epsilon(\mathbb{R}^\nu)$, where $\Lambda_\epsilon: \mathbb{R}^\nu \rightarrow H^1(\Omega)$ is a map obtained from the contraction mapping principle applied to a properly defined nonlinear operator Γ_ϵ defined on a certain space of maps $y:]-\infty, 0] \rightarrow H^1(\Omega)$ of exponential growth. If the operator Γ_ϵ is a contraction then the map Λ_ϵ is well-defined and $\mathcal{A}_\epsilon \subset \mathcal{M}_\epsilon$. It follows that $\mathcal{M}_\epsilon$ is invariant with respect to solutions of the original semiflow $\pi_{\epsilon, \hat{f}}$ as long as these solutions stay in the open set U . One can even find an open set $V \subset \mathbb{R}^\nu$ such that for $\epsilon \in [0, \epsilon_0]$ the set $\Lambda_\epsilon(V)$ is positively invariant with respect to $\pi_{\epsilon, \hat{f}}$ and $\mathcal{A}_\epsilon \subset \Lambda_\epsilon(V) \subset U$.

The only problem is that, under the usual norm $|\cdot|_\epsilon$ on $H^1(\Omega)$, the operator Γ_ϵ is **not** a contraction. At this point we use an ingenious idea due to Brunovský and Tereščák (see Theorem 4.1 in [3] and its proof) and, given positive numbers l and L introduce an equivalent norm

$$\|u\|_\epsilon = L|u|_{L^2} + l|u|_\epsilon$$

on $H^1(\Omega)$. Similarly as in [3] we seek to choose the constants l and L in such a way that the operator Γ_ϵ is a uniform contraction with respect to the norm $\|\cdot\|_\epsilon$. That this is possible follows, on the one hand, from the linear estimates contained in Lemma 5.3 below and, on the other hand, from the following C^1 -cut-off-result for Nemitski operators:

Theorem 5.1. *Let Ω be an open, bounded subset of $\mathbb{R}^2$, with Lipschitz boundary. Let $f \in C^1(\mathbb{R} \rightarrow \mathbb{R})$ satisfy the growth estimate*

$$|f'(s)| \leq C(1 + |s|^\beta) \quad \text{for } s \in \mathbb{R}$$

where C and $\beta \in [0, \infty[$ are arbitrary real constants.

Let l be an arbitrary positive real number and B be an arbitrary bounded subset of $H^1(\Omega)$. Then there exists an open set $U = U(l, B) \subset H^1(\Omega)$ including B , a positive real number $L = L(l, B)$ and a map $g = g(l, B) \in C^1(H^1(\Omega) \rightarrow L^2(\Omega))$ with $\hat{f}(u) = g(u)$ for $u \in U$ and such that g maps $H_s^1(\Omega)$ into $L_s^2(\Omega)$ and satisfies the estimates

$$(5.1) \quad \sup_{u \in H^1(\Omega)} |g(u)|_{L^2} < \infty$$

$$(5.2) \quad |g(u) - g(v)|_{L^2} \leq L|u - v|_{L^2} + l|u - v|_{H^1} \quad \text{for } u, v \in H^1(\Omega)$$

and

$$(5.3) \quad |Dg(u)v|_{L^2} \leq L|v|_{L^2} + l|v|_{H^1} \quad \text{for } u, v \in H^1(\Omega).$$

Proof. We can assume, without loss of generality, that $\beta > 0$. Take a real number $q > 2$ (so in particular $0 < 1 - 2/q < 1$ and $q \geq 2(\beta + 1)$). Choose θ with $1 - 2/q < \theta < 1$. By the Gagliardo-Nirenberg inequality there is a constant C_1 such that

$$|u|_{L^q} \leq C_1 |u|_{H^1}^\theta |u|_{L^2}^{1-\theta} \quad \text{for } u \in H^1(\Omega).$$

Hence, for every $\rho > 0$

$$(5.4) \quad |u|_{L^q} \leq C_1 \theta \rho^{1-\theta} |u|_{H^1} + C_1 (1 - \theta) \rho^{-\theta} |u|_{L^2} \quad \text{for } u \in H^1(\Omega).$$

In particular, this implies that there is a real positive constant M with

$$|u|_{L^q} < M \quad \text{for } u \in B.$$

Define U to be the set of all $u \in H^1(\Omega)$ with $|u|_{L^q} < M$. It follows that U is open in $H^1(\Omega)$ and $B \subset U$. We now apply Theorem 2.1 and obtain that $\hat{f}$ is a C^1 -map from $L^q(\Omega)$ to $L^2(\Omega)$ such that both $|\hat{f}(u)|_{L^2}$ and $|D\hat{f}(u)|_{L(L^q(\Omega) \rightarrow L^2(\Omega))}$ are bounded on bounded subsets of $L^q(\Omega)$. It is well-known that there is a bounded function $h \in C^1(L^q(\Omega) \rightarrow \mathbb{R})$ having globally bounded Fréchet derivative and such that $h(u) = 1$ if $|u|_{L^q} \leq M$ and $h(u) = 0$ if $|u|_{L^q} > 2M$. This can also easily be proved using Theorem 2.1. In fact, since the function $s \mapsto |s|^q$ is in $C^1(\mathbb{R} \rightarrow \mathbb{R})$ with derivative $s \mapsto q|s|^{q-1}$, Theorem 2.1 implies that the map $\xi: u \mapsto |u|^q$ is in $C^1(L^q(\Omega) \rightarrow L^1(\Omega))$. Furthermore, on bounded sets in $L^q(\Omega)$ the map ξ is bounded and has bounded Fréchet derivative. Let $\phi \in C^1(\mathbb{R} \rightarrow \mathbb{R})$ be such that $\phi(x) = 1$ if $|x| \leq M^q$ and $\phi(x) = 0$ if $|x| > (2M)^q$. It follows that the function $h(u) := \phi(|u|_{L^q}^q)$ for $u \in L^q(\Omega)$ has the desired properties. Define $g(u) = h(u)\hat{f}(u)$ for $u \in L^q(\Omega)$. It follows using Leibniz rule that g is a C^1 -map from $L^q(\Omega)$ to $L^2(\Omega)$ such that both $|g(u)|_{L^2}$ and $|Dg(u)|_{L(L^q(\Omega) \rightarrow L^2(\Omega))}$ are globally bounded. It follows that g is globally Lipschitzian from $L^q(\Omega)$ to $L^2(\Omega)$ with a Lipschitz constant K . Using (5.4) it follows that for every $\rho > 0$ and all $u, v \in H^1(\Omega)$

$$|g(u) - g(v)|_{L^2} \leq C_1 K (1 - \theta) \rho^{-\theta} |u - v|_{L^2} + C_1 K \theta \rho^{1-\theta} |u - v|_{H^1}.$$

Choose $\rho > 0$ so small that $C_1 K \theta \rho^{1-\theta} \leq l$ and set $L = C_1 K (1 - \theta) \rho^{-\theta}$. Since $H^1(\Omega)$ is continuously contained in $L^q(\Omega)$ it follows that g is of class C^1 as a map from $H^1(\Omega)$ to $L^2(\Omega)$ and that (5.1) and (5.2) are satisfied. Moreover, g maps $H_s^1(\Omega)$ into $L_s^2(\Omega)$ since $\hat{f}$ does so. The estimate (5.2) easily implies (5.3). Finally, the definition of U immediately implies that $\hat{f}(u) = g(u)$ for $u \in U$. The theorem is proved. $\square$

In order to state our main result, we need some notation:

For every $\epsilon \in [0, 1]$ and every $\nu \in \mathbb{N}$ let $X_{\epsilon, \nu, 1}$ be the span of the first ν eigenvectors $w_{\epsilon, j}$, $j = 1, \dots, \nu$, of A_ϵ . Let $X_{\epsilon, \nu, 2}$ be the orthogonal complement of $X_{\epsilon, \nu, 1}$ in $L^2(\Omega)$ if $\epsilon > 0$ and in $L_s^2(\Omega)$ if $\epsilon = 0$. Let $A_{\epsilon, \nu, i}$ be the restriction of A_ϵ to $X_{\epsilon, \nu, i}$ for $i = 1, 2$. Let $E_{\epsilon, \nu} \xi := \sum_{j=1}^{\nu} \xi_j w_{\epsilon, j}$, $\xi \in \mathbb{R}^\nu$ and $P_{\epsilon, \nu, i}$ be the orthogonal projection of $L^2(\Omega)$ onto $X_{\epsilon, \nu, i}$, $i = 1, 2$ if $\epsilon > 0$ and $P_{0, \nu, i}$ be the orthogonal projection of $L_s^2(\Omega)$ onto $X_{0, \nu, i}$, $i = 1, 2$.

Theorem 5.2. *Suppose that $f \in C^1(\mathbb{R} \rightarrow \mathbb{R})$ is dissipative in the sense that*

$$\limsup_{|s| \rightarrow \infty} f(s)/s \leq -\delta_0 \quad \text{for some } \delta_0 > 0.$$

Furthermore, let f satisfy the growth estimate

$$|f'(s)| \leq C(1 + |s|^\beta) \quad \text{for } s \in \mathbb{R},$$

where C and $\beta \in [0, \infty[$ are arbitrary real constants. Suppose the eigenvalues of A_0 satisfy the following gap condition:

$$(5.5) \quad \limsup_{\nu \rightarrow \infty} \frac{\lambda_{0, \nu+1} - \lambda_{0, \nu}}{\lambda_{0, \nu}^{1/2}} > 0.$$

Then there are an $\epsilon_0 > 0$ and an open bounded set $U \subset H^1(\Omega)$ such that for every $\epsilon \in [0, \epsilon_0[$ the attractor $\mathcal{A}_\epsilon$ of the semiflow $\pi_{\epsilon, \hat{f}}$ lies in U .

Furthermore, there exists a globally Lipschitzian map $g \in C^1(H^1(\Omega) \rightarrow L^2(\Omega))$ with $g(u) = \hat{f}(u)$ for $u \in U$.

Besides, there is a positive integer ν and for every $\epsilon \in [0, \epsilon_0[$ there is a map $\Lambda_\epsilon \in C^1(\mathbb{R}^\nu \rightarrow H^1(\Omega))$ if $\epsilon > 0$ and $\Lambda_0 \in C^1(\mathbb{R}^\nu \rightarrow H_s^1(\Omega))$ such that

$$(5.6) \quad P_{\epsilon, \nu, 1} \circ \Lambda_\epsilon = E_{\epsilon, \nu}$$

and $\Lambda_\epsilon(\mathbb{R}^\nu)$ is an invariant manifold with respect to the semiflow $\pi_{\epsilon, g}$.

Finally, there is an open set $V \subset \mathbb{R}^\nu$ such that for every $\epsilon \in [0, \epsilon_0[$

$$\mathcal{A}_\epsilon \subset \Lambda_\epsilon(V) \subset U$$

and the set $\Lambda_\epsilon(V)$ is positively invariant with respect to the semiflow $\pi_{\epsilon, \hat{f}}$.

The reduced equation on $\Lambda_\epsilon(\mathbb{R}^\nu)$ takes the form

$$(5.7) \quad \dot{\xi} = v_\epsilon(\xi), \quad \xi \in \mathbb{R}^\nu,$$

where

$$v_\epsilon: \mathbb{R}^\nu \rightarrow \mathbb{R}^\nu, \quad \xi \mapsto -A_\epsilon E_{\epsilon, \nu} \xi + P_{\epsilon, \nu, 1} g(\Lambda_\epsilon(\xi)).$$

Moreover, whenever $\epsilon_n \rightarrow 0^+$ and $\xi_n \rightarrow \xi_0$ in $\mathbb{R}^\nu$, then

$$(5.8) \quad |\Lambda_{\epsilon_n}(\xi_n) - \Lambda_0(\xi_0)|_{\epsilon_n} + \sum_{j=1}^{\nu} |\partial_j \Lambda_{\epsilon_n}(\xi_n) - \partial_j \Lambda_0(\xi_0)|_{\epsilon_n} \rightarrow 0$$

and

$$(5.9) \quad |v_{\epsilon_n}(\xi_n) - v_0(\xi_0)|_{\mathbb{R}^\nu} + \sum_{j=1}^{\nu} |\partial_j v_{\epsilon_n}(\xi_n) - \partial_j v_0(\xi_0)|_{\mathbb{R}^\nu} \rightarrow 0.$$

Given $\mu \in \mathbb{R}$, a Banach space $(Y, |\cdot|_Y)$ and a function $y:]-\infty, 0] \rightarrow Y$ we write

$$|y|_{\mu, |\cdot|_Y} := \sup_{t \in]-\infty, 0]} e^{\mu t} |y(t)|_Y$$

and we denote by $BC^\mu(Y, |\cdot|_Y)$ the set of all continuous functions $y:]-\infty, 0] \rightarrow Y$ such that $|y|_{\mu, |\cdot|_Y} < \infty$. The space $BC^\mu(Y, |\cdot|_Y)$ is a Banach space with respect to the norm $y \mapsto |y|_{\mu, |\cdot|_Y}$. In particular, we write

- (1) $BC^\mu(L^2(\Omega)) := BC^\mu(L^2(\Omega), |\cdot|_{L^2})$, with the norm $|y|_{\mu, L^2} := |y|_{\mu, |\cdot|_{L^2}}$,
- (2) $BC^\mu(L_s^2(\Omega)) := BC^\mu(L_s^2(\Omega), |\cdot|_{L^2})$, with the norm $|y|_{\mu, L^2} := |y|_{\mu, |\cdot|_{L^2}}$,
- (3) $BC^\mu(H^1(\Omega), \epsilon) := BC^\mu(H^1(\Omega), |\cdot|_\epsilon)$ with the norm $|y|_{\mu, \epsilon} := |y|_{\mu, |\cdot|_\epsilon}$ for $\epsilon > 0$,
- (4) $BC^\mu(H^1(\Omega), 0) := BC^\mu(H_s^1(\Omega), |\cdot|_{H^1})$ with the norm $|y|_{\mu, 0} := |y|_{\mu, |\cdot|_{H^1}}$.

We need the following Lemma:

Lemma 5.3.

- (1) For every $\epsilon \in [0, 1]$ and every $\nu \in \mathbb{N}$

$$|e^{-A_{\epsilon, \nu, 1} t} u|_{L^2} \leq e^{-\lambda_{\epsilon, \nu} t} |u|_{L^2} \quad u \in X_{\epsilon, \nu, 1}, t \leq 0,$$

$$|e^{-A_{\epsilon, \nu, 2} t} u|_{L^2} \leq e^{-\lambda_{\epsilon, \nu+1} t} |u|_{L^2} \quad u \in X_{\epsilon, \nu, 2}, t > 0,$$

$$|e^{-A_{\epsilon, \nu, 1} t} u|_\epsilon \leq (\lambda_{\epsilon, \nu} + 1)^{1/2} e^{-\lambda_{\epsilon, \nu} t} |u|_{L^2} \quad u \in X_{\epsilon, \nu, 1}, t \leq 0,$$

$$|e^{-A_{\epsilon, \nu, 2} t} u|_\epsilon \leq \left((\lambda_{\epsilon, \nu+1} + 1)^{1/2} + C_{1/2} t^{-1/2} \right) e^{-\lambda_{\epsilon, \nu+1} t} |u|_{L^2} \quad u \in X_{\epsilon, \nu, 2}, t > 0.$$

- (2) Define the operator

$$\Xi_{\epsilon, \nu}(\xi, y)(t) = e^{-A_{\epsilon, \nu, 1} t} E_{\epsilon, \nu} \xi + K_{\epsilon, \nu} y(t)$$

where $\xi \in \mathbb{R}^\nu$, and for $y:]-\infty, 0] \rightarrow L^2(\Omega)$ and $t \leq 0$

$$(5.10) \quad K_{\epsilon, \nu} y(t) = \int_0^t e^{-A_{\epsilon, \nu, 1}(t-s)} P_{\epsilon, \nu, 1} y(s) ds + \int_{-\infty}^t e^{-A_{\epsilon, \nu, 2}(t-s)} P_{\epsilon, \nu, 2} y(s) ds,$$

whenever the right hand side of (5.10) makes sense. Let ζ with $\lambda_{\epsilon,\nu} < \zeta < \lambda_{\epsilon,\nu+1}$ be arbitrary. Then $\Xi_{\epsilon,\nu}$ maps $\mathbb{R}^\nu \times BC^\zeta(L^2(\Omega))$ into $BC^\zeta(H^1(\Omega), \epsilon)$ for $\epsilon > 0$ and $\Xi_{0,\nu}$ maps $\mathbb{R}^\nu \times BC^\zeta(L_s^2(\Omega))$ into $BC^\zeta(H^1(\Omega), 0)$. Moreover, for $\epsilon > 0$ and $y \in BC^\zeta(L^2(\Omega))$ (resp. for $\epsilon = 0$ and $y \in BC^\zeta(L_s^2(\Omega))$)

$$(5.11) \quad |K_{\epsilon,\nu}y|_{\zeta,L^2} \leq \left(\frac{1}{\zeta - \lambda_{\epsilon,\nu}} + \frac{1}{\lambda_{\epsilon,\nu+1} - \zeta} \right) |y|_{\zeta,L^2},$$

and

$$(5.12) \quad |K_{\epsilon,\nu}y|_{\zeta,\epsilon} \leq \left(\frac{(\lambda_{\epsilon,\nu} + 1)^{1/2}}{\zeta - \lambda_{\epsilon,\nu}} + \frac{(\lambda_{\epsilon,\nu+1} + 1)^{1/2}}{\lambda_{\epsilon,\nu+1} - \zeta} + C'_{1/2}(\lambda_{\epsilon,\nu+1} - \zeta)^{-1/2} \right) |y|_{\zeta,L^2}.$$

(3) If $\lambda_{0,\nu} < \zeta < \lambda_{0,\nu+1}$, $\epsilon_n \rightarrow 0^+$, $\xi_n \rightarrow \xi_0$ in $\mathbb{R}^\nu$ and $y_n \rightarrow y_0$ in $BC^\zeta(L^2(\Omega))$, where $y_0 \in BC^\zeta(L_s^2(\Omega))$, then, for all n large enough, $\lambda_{\epsilon_n,\nu} < \zeta < \lambda_{\epsilon_n,\nu+1}$ and

$$|\Xi_{\epsilon_n,\nu}(\xi_n, y_n) - \Xi_{0,\nu}(\xi_0, y_0)|_{\zeta,\epsilon_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Proof of the lemma. The proof of part (1) is obvious and well-known. Part (2) follows from part (1) and well-known results about semigroups generated by sectorial operators. We prove (5.12) for $\epsilon > 0$, the other cases being completely analogous. Let $y \in BC^\zeta(L^2(\Omega))$ and $t \leq 0$. By the definition of $K_{\epsilon,\nu}$ and by the estimates of part (1), we have

$$\begin{aligned} |K_{\epsilon,\nu}y(t)|_\epsilon &\leq (\lambda_{\epsilon,\nu} + 1)^{1/2} e^{-\lambda_{\epsilon,\nu}t} |y|_{\zeta,L^2} \left| \int_0^t e^{(\lambda_{\epsilon,\nu} - \zeta)s} ds \right| \\ &\quad + (\lambda_{\epsilon,\nu+1} + 1)^{1/2} e^{-\lambda_{\epsilon,\nu+1}t} |y|_{\zeta,L^2} \int_{-\infty}^t e^{(\lambda_{\epsilon,\nu+1} - \zeta)s} ds \\ &\quad + C_{1/2} e^{-\zeta t} |y|_{\zeta,L^2} \int_{-\infty}^t (t-s)^{-1/2} e^{-(\lambda_{\epsilon,\nu+1} - \zeta)(t-s)} ds. \end{aligned}$$

Then (5.12) follows from simple integration.

Let us prove part (3). Suppose that $\lambda_{0,\nu} < \zeta < \lambda_{0,\nu+1}$, $\epsilon_n \rightarrow 0^+$, $\xi_n \rightarrow \xi_0$ in $\mathbb{R}^\nu$ and $y_n \rightarrow y_0$ in $BC^\zeta(L^2(\Omega))$. Since $\lambda_{\epsilon,\nu} \rightarrow \lambda_{0,\nu}$ and $\lambda_{\epsilon,\nu+1} \rightarrow \lambda_{0,\nu+1}$ as $\epsilon \rightarrow 0^+$ (see [15, Th. 3.3]), it follows that there is an $n_0 \in \mathbb{N}$ such that $\lambda_{\epsilon_n,\nu} < \zeta < \lambda_{\epsilon_n,\nu+1}$ for all $n \geq n_0$. In the rest of this proof let $n \geq n_0$. Set $z_n(t) := e^{\zeta t} y_n(t)$ and $z_0(t) := e^{\zeta t} y_0(t)$ for $t \leq 0$. It easily follows that

$$\sup_{r \in]-\infty, 0]} |z_n(r) - z_0(r)|_{L^2} \rightarrow 0.$$

Thus, by Theorem 3.3 in [15], for every $j \in \mathbb{N}$

$$\sup_{r \in]-\infty, 0]} |\langle z_n(r), w_{\epsilon_n,j} \rangle w_{\epsilon_n,j} - \langle z_0(r), w_{0,j} \rangle w_{0,j}|_{L^2}$$

$$\begin{aligned} &\leq \sup_{r \in]-\infty, 0]} |z_n(r) - z_0(r)|_{L^2} \\ &+ 2 \sup_{r \in]-\infty, 0]} |z_0(r)|_{L^2} |w_{\epsilon_n, j} - w_{0, j}|_{L^2} \rightarrow 0. \end{aligned}$$

Therefore

$$(5.13) \quad \sup_{r \in]-\infty, 0]} |P_{\epsilon_n, \nu, 1} z_n(r) - P_{0, \nu, 1} z_0(r)|_{L^2} \rightarrow 0$$

and so

$$(5.14) \quad \sup_{r \in]-\infty, 0]} |P_{\epsilon_n, \nu, 2} z_n(r) - P_{0, \nu, 2} z_0(r)|_{L^2} \rightarrow 0.$$

Note that whenever (a_n) is a sequence of positive numbers converging to some positive number a , then

$$(5.15) \quad \sup_{r \in]-\infty, 0]} |e^{a_n t} - e^{at}| \rightarrow 0.$$

Consequently, again by Theorem 3.3 in [15],

$$\begin{aligned} &\sup_{t \in]-\infty, 0]} e^{\zeta t} |e^{-A_{\epsilon_n, \nu, 1} t} E_{\epsilon_n, \nu} \xi_n - e^{-A_{0, \nu, 1} t} E_{0, \nu} \xi_0|_{\epsilon_n} \\ &\leq \sum_{j=1}^{\nu} \sup_{t \in]-\infty, 0]} \left| e^{(\zeta - \lambda_{\epsilon_n, j})t} \xi_{n, j} w_{\epsilon_n, j} - e^{(\zeta - \lambda_{0, j})t} \xi_{0, j} w_{0, j} \right|_{\epsilon_n} \\ &\leq \sup_{t \in]-\infty, 0]} \left| e^{(\zeta - \lambda_{\epsilon_n, j})t} - e^{(\zeta - \lambda_{0, j})t} \right| |\xi_{0, j} w_{0, j}|_{\epsilon_n} \\ &+ \sup_{t \in]-\infty, 0]} e^{(\zeta - \lambda_{0, j})t} |\xi_{n, j} w_{\epsilon_n, j} - \xi_{0, j} w_{0, j}|_{\epsilon_n} \rightarrow 0. \end{aligned}$$

Furthermore,

$$\begin{aligned} &\sup_{t \leq 0} e^{\zeta t} \left| \int_0^t e^{-A_{\epsilon_n, \nu, 1}(t-s)} P_{\epsilon_n, \nu, 1} y_n(s) ds - \int_0^t e^{-A_{0, \nu, 1}(t-s)} P_{0, \nu, 1} y_0(s) ds \right|_{\epsilon_n} \leq \\ &\sum_{j=1}^{\nu} \sup_{t \leq 0} \int_t^0 \left| e^{(\zeta - \lambda_{\epsilon_n, j})(t-s)} \langle z_n(s), w_{\epsilon_n, j} \rangle w_{\epsilon_n, j} - e^{(\zeta - \lambda_{0, j})(t-s)} \langle z_0(s), w_{0, j} \rangle w_{0, j} \right|_{\epsilon_n} ds \\ &\leq \sum_{j=1}^{\nu} \sup_{t \leq 0} \int_t^0 \left| e^{(\zeta - \lambda_{\epsilon_n, j})(t-s)} - e^{(\zeta - \lambda_{0, j})(t-s)} \right| |\langle z_n(s), w_{\epsilon_n, j} \rangle w_{\epsilon_n, j}|_{\epsilon_n} ds \\ &+ \sum_{j=1}^{\nu} \sup_{t \leq 0} \int_t^0 e^{(\zeta - \lambda_{0, j})(t-s)} |\langle z_n(s), w_{\epsilon_n, j} \rangle w_{\epsilon_n, j} - \langle z_0(s), w_{0, j} \rangle w_{0, j}|_{\epsilon_n} ds \end{aligned}$$

$$\begin{aligned}
&\leq \sum_{j=1}^{\nu} \sup_{t \leq 0} \int_t^0 \left| e^{(\zeta - \lambda_{\epsilon_n, j})(t-s)} - e^{(\zeta - \lambda_{0, j})(t-s)} \right| ds \cdot \sup_{r \leq 0} |\langle z_n(r), w_{\epsilon_n, j} \rangle w_{\epsilon_n, j}|_{\epsilon_n} \\
&+ \sum_{j=1}^{\nu} \sup_{t \leq 0} \int_t^0 e^{(\zeta - \lambda_{0, j})(t-s)} ds \cdot \sup_{r \leq 0} |\langle z_n(r), w_{\epsilon_n, j} \rangle w_{\epsilon_n, j} - \langle z_0(r), w_{0, j} \rangle w_{0, j}|_{\epsilon_n} =: T_n.
\end{aligned}$$

Since

$$\begin{aligned}
\int_t^0 \left| e^{(\zeta - \lambda_{\epsilon_n, j})(t-s)} - e^{(\zeta - \lambda_{0, j})(t-s)} \right| ds &= \left| \int_t^0 e^{(\zeta - \lambda_{\epsilon_n, j})(t-s)} - e^{(\zeta - \lambda_{0, j})(t-s)} ds \right| \\
&= \left| \frac{1}{\zeta - \lambda_{\epsilon_n, j}} (1 - e^{(\zeta - \lambda_{\epsilon_n, j})(t-s)}) - \frac{1}{\zeta - \lambda_{0, j}} (1 - e^{(\zeta - \lambda_{0, j})(t-s)}) \right|
\end{aligned}$$

we obtain from (5.15) and Theorem 3.3 in [15] that

$$T_n \rightarrow 0.$$

We also have

$$\begin{aligned}
&\sup_{t \leq 0} e^{\zeta t} \left| \int_{-\infty}^t e^{-A_{\epsilon_n, \nu, 2}(t-s)} P_{\epsilon_n, \nu, 2} y_n(s) ds - \int_{-\infty}^t e^{-A_{0, \nu, 2}(t-s)} P_{0, \nu, 2} y_0(s) ds \right|_{\epsilon_n} \\
&\leq \sup_{t \leq 0} \int_{-\infty}^t \left| e^{-A_{\epsilon_n, \nu, 2}(t-s)} e^{\zeta(t-s)} P_{\epsilon_n, \nu, 2} z_n(s) - e^{-A_{0, \nu, 2}(t-s)} e^{\zeta(t-s)} P_{0, \nu, 2} z_0(s) \right|_{\epsilon_n} ds \\
&= \sup_{t \leq 0} \int_0^{\infty} \left| e^{-A_{\epsilon_n, \nu, 2}s} e^{\zeta s} P_{\epsilon_n, \nu, 2} z_n(t-s) - e^{-A_{0, \nu, 2}s} e^{\zeta s} P_{0, \nu, 2} z_0(t-s) \right|_{\epsilon_n} ds.
\end{aligned}$$

Part (1) of this lemma implies that there is an integrable function $h: [0, \infty[\rightarrow [0, \infty[$ such that

$$\left| e^{-A_{\epsilon_n, \nu, 2}s} e^{\zeta s} P_{\epsilon_n, \nu, 2} z_n(t-s) - e^{-A_{0, \nu, 2}s} e^{\zeta s} P_{0, \nu, 2} z_0(t-s) \right|_{\epsilon_n} \leq h(s)$$

for all $n \in \mathbb{N}$, and $t, s \leq 0$.

Therefore the dominated convergence theorem, together with (5.14) and Theorem 4.1 in [15] imply that

$$\sup_{t \leq 0} \int_0^{\infty} \left| e^{-A_{\epsilon_n, \nu, 2}s} e^{\zeta s} P_{\epsilon_n, \nu, 2} z_n(t-s) - e^{-A_{0, \nu, 2}s} e^{\zeta s} P_{0, \nu, 2} z_0(t-s) \right|_{\epsilon_n} ds \rightarrow 0.$$

Putting everything together we obtain

$$|\Xi_{\epsilon_n, \nu}(\xi_n, y_n) - \Xi_{0, \nu}(\xi_0, y_0)|_{\zeta, \epsilon_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

The lemma is proved. $\square$

Proof of Theorem 5.2. We divide the proof into several steps:

Step 1: choice of l and U .

For $\nu \in \mathbb{N}$ with $\lambda_{0,\nu+1} - \lambda_{0,\nu} > 0$ define

$$\eta_\nu = (\lambda_{0,\nu+1} - \lambda_{0,\nu})/5$$

and

$$I_\nu = [\lambda_{0,\nu} + 2\eta_\nu, \lambda_{0,\nu} + 3\eta_\nu].$$

It follows that

$$\zeta - \lambda_{0,\nu} > (\lambda_{0,\nu+1} - \lambda_{0,\nu})/3 \text{ and } \lambda_{0,\nu+1} - \zeta > (\lambda_{0,\nu+1} - \lambda_{0,\nu})/3 \text{ for } \zeta \in I_\nu.$$

Hence

$$(5.16) \quad \sup_{\zeta \in I_\nu} \left(\frac{1}{\zeta - \lambda_{0,\nu}} + \frac{1}{\lambda_{0,\nu+1} - \zeta} \right) < C_{\nu,1} := 6 \frac{1}{\lambda_{0,\nu+1} - \lambda_{0,\nu}}$$

and

$$(5.17) \quad \sup_{\zeta \in I_\nu} \left(\frac{(\lambda_{0,\nu} + 1)^{1/2}}{\zeta - \lambda_{0,\nu}} + \frac{(\lambda_{0,\nu+1} + 1)^{1/2}}{\lambda_{0,\nu+1} - \zeta} + C'_{1/2}(\lambda_{0,\nu+1} - \zeta)^{-1/2} \right) < C_{\nu,2},$$

where

$$C_{\nu,2} := 3 \frac{(\lambda_{0,\nu} + 1)^{1/2}}{\lambda_{0,\nu+1} - \lambda_{0,\nu}} + 3 \frac{(\lambda_{0,\nu+1} + 1)^{1/2}}{\lambda_{0,\nu+1} - \lambda_{0,\nu}} + 3^{1/2} C'_{1/2}(\lambda_{0,\nu+1} - \lambda_{0,\nu})^{-1/2}.$$

In view of (5.5) there is a $C_1 \in [0, \infty[$ and a strictly increasing sequence $(\nu_k)_{k \in \mathbb{N}}$ in $\mathbb{N}$ such that $\lambda_{0,\nu_k+1} - \lambda_{0,\nu_k} > 0$ for $k \in \mathbb{N}$ and

$$\lambda_{0,\nu_k}^{1/2} / (\lambda_{0,\nu_k+1} - \lambda_{0,\nu_k}) \rightarrow C_1 \quad \text{as } k \rightarrow \infty.$$

Since $\lambda_{0,\nu} \rightarrow \infty$ as $\nu \rightarrow \infty$ it follows that

$$\frac{1}{\lambda_{0,\nu_k+1} - \lambda_{0,\nu_k}} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Consequently,

$$(5.18) \quad C_{\nu_k,1} \rightarrow 0 \quad \text{and } C_{\nu_k,2} \rightarrow 6C_1 \text{ as } k \rightarrow \infty.$$

Choose l such that

$$0 < l(6C_1 + 1) < \frac{1}{4}.$$

By Theorem 5.10 in [15] there is an ϵ_0 with $0 < \epsilon_0 < 1$ and a bounded set B_1 in $H^1(\Omega)$ such that for every $\epsilon \in [0, \epsilon_0]$ the attractor $\mathcal{A}_\epsilon$ lies in B_1 . Let V_0 be the Liapunov function of $\pi_{0,\hat{f}}$ defined by

$$V_0: H_s^1(\Omega) \rightarrow \mathbb{R}, \quad u \mapsto (1/2)|\nabla u|_{L^2}^2 - \int_{\Omega} \hat{F}(u) \, dx dy.$$

Here, as usual, $\hat{F}: H^1(\Omega) \rightarrow L^1(\Omega)$ is the Nemitski operator defined by the function $F(x) := \int_0^x f(s) \, ds$, $x \in \mathbb{R}$. There is an $M_0 \in]0, \infty[$ so that $V_0(u) < M_0$ for all $u \in \mathcal{A}_0$. As in [15] it follows that

$$B_2 := \{ u \in H_s^1(\Omega) \mid V_0(u) \leq M_0 \}$$

is bounded. Define $B := B_1 \cup B_2$ and let $L = L(l, B)$ and $U = U(l, B)$ and $g = g(l, B)$ be as in Corollary 5.1. By (5.18) there exists a $k \in \mathbb{N}$ such that

$$(5.19) \quad LC_{\nu_k,1} < \frac{1}{4} \quad \text{and} \quad lC_{\nu_k,2} < \frac{1}{4}.$$

Fix such a k and set $\nu := \nu_k$. Since $\lambda_{\epsilon,\nu} \rightarrow \lambda_{0,\nu}$ and $\lambda_{\epsilon,\nu+1} \rightarrow \lambda_{0,\nu+1}$ as $\epsilon \rightarrow 0^+$ and using (5.16) and (5.17), we may assume, by taking ϵ_0 smaller if necessary, that for every $\epsilon \in [0, \epsilon_0]$ and every $\zeta \in I_\nu$

$$\lambda_{\epsilon,\nu} < \zeta < \lambda_{\epsilon,\nu+1},$$

$$(5.20) \quad \sup_{\zeta \in I_\nu} \left(\frac{1}{\zeta - \lambda_{\epsilon,\nu}} + \frac{1}{\lambda_{\epsilon,\nu+1} - \zeta} \right) < C_{\nu,1}$$

and

$$(5.21) \quad \sup_{\zeta \in I_\nu} \left(\frac{(\lambda_{\epsilon,\nu} + 1)^{1/2}}{\zeta - \lambda_{\epsilon,\nu}} + \frac{(\lambda_{\epsilon,\nu+1} + 1)^{1/2}}{\lambda_{\epsilon,\nu+1} - \zeta} + C'_{1/2}(\lambda_{\epsilon,\nu+1} - \zeta)^{-1/2} \right) < C_{\nu,2}.$$

Step 2: choice of a new norm.

If $\epsilon > 0$ endow $H^1(\Omega)$ with the equivalent norm

$$\|u\|_\epsilon := L|u|_{L^2} + l|u|_\epsilon.$$

Write $Z_\epsilon^\zeta := BC^\zeta(H^1(\Omega), \|\cdot\|_\epsilon)$ with the corresponding norm

$$\|y\|_{\zeta,\epsilon} = \sup_{t \in]-\infty, 0]} e^{\zeta t} \|y(t)\|_\epsilon.$$

If $\epsilon = 0$ endow $H_s^1(\Omega)$ with the equivalent norm

$$\|u\|_0 := L|u|_{L^2} + l|u|_{H^1}.$$

Write $Z_0^\zeta := BC^\zeta(H_s^1(\Omega), \|\cdot\|_0)$ with the corresponding norm

$$\|y\|_{\zeta,0} = \sup_{t \in]-\infty, 0]} e^{\zeta t} \|y(t)\|_0.$$

It follows that for $\epsilon \in [0, \epsilon_0]$, $\zeta \in I_\nu$ and $y \in BC^\zeta(L^2(\Omega))$ (resp. for $\epsilon = 0$ and $y \in BC^\zeta(L_s^2(\Omega))$)

$$(5.22) \quad \|K_{\epsilon,\nu} y\|_{\zeta,\epsilon} \leq (1/2) |y|_{\zeta, L^2}.$$

Step 3: the contraction map Γ_ϵ .

Fix ζ and $\mu \in I_\nu$ with $\zeta < \mu$. Since $g: H^1(\Omega) \rightarrow L^2(\Omega)$ is globally bounded, it follows that $g \circ y$ is globally bounded for every $y \in BC^\zeta(H^1(\Omega), \epsilon)$, $\epsilon \geq 0$. Moreover, since g maps $H_s^1(\Omega)$ into $L_s^2(\Omega)$ it follows that the nonlinear operator $y \mapsto g \circ y$ maps Z_ϵ^ζ into $BC^\zeta(L^2(\Omega))$ for $\epsilon > 0$ and it maps Z_0^ζ into $BC^\zeta(L_s^2(\Omega))$. Moreover,

$$\begin{aligned} |g(u) - g(v)|_{L^2} &\leq L|u - v|_{L^2} + l|u - v|_{H^1} \\ &\leq \|u - v\|_\epsilon \quad \text{for } u, v \in H^1(\Omega) \end{aligned}$$

so

$$(5.23) \quad |g \circ y - g \circ w|_{\zeta, L^2} \leq \|y - w\|_{\zeta, \epsilon} \quad \text{for } y, w \in Z_\epsilon^\zeta.$$

It follows that the operator

$$\Gamma_\epsilon: \mathbb{R}^\nu \times Z_\epsilon^\zeta \rightarrow Z_\epsilon^\zeta, \quad (\xi, y) \mapsto \Xi_{\epsilon,\nu}(\xi, g \circ y)$$

is well-defined. Spelled out in full, the operator Γ_ϵ reads as follows:

$$\begin{aligned} \Gamma_\epsilon(\xi, y)(t) &= e^{-A_{\epsilon,\nu,1}t} E_{\epsilon,\nu} \xi + \int_0^t e^{-A_{\epsilon,\nu,1}(t-s)} P_{\epsilon,\nu,1} g(y(s)) \, ds \\ &+ \int_{-\infty}^t e^{-A_{\epsilon,\nu,2}(t-s)} P_{\epsilon,\nu,2} g(y(s)) \, ds \quad \text{for } (\xi, y) \in \mathbb{R}^\nu \times Z_\epsilon^\zeta \text{ and } t \leq 0. \end{aligned}$$

If $0 \leq \epsilon \leq \epsilon_0$ then, by (5.22) and (5.23), Γ_ϵ is a uniform contraction in the second variable with contraction constant $1/2$. It follows that for every such ϵ there is a uniquely defined map $\phi_\epsilon: \mathbb{R}^\nu \rightarrow Z_\epsilon^\zeta$, such that

$$\phi_\epsilon(\xi) = \Gamma_\epsilon(\xi, \phi_\epsilon(\xi)) \quad \text{for } \xi \in \mathbb{R}^\nu.$$

Either proceeding directly, or by using the fiber contraction theorem as in [4] or else by using the abstract results of [18] one proves that the map ϕ_ϵ is of class C^1

as a map from $\mathbb{R}^\nu$ into Z_ϵ^μ and for every $\xi' \in \mathbb{R}^\nu$ and $j = 1, \dots, \nu$, $D\phi_\epsilon(\xi)\xi'$ lies in Z_ϵ^ζ and is given by the recursive formula

$$(5.24) \quad D\phi_\epsilon(\xi)\xi' = \Xi_{\epsilon,\nu}(\xi', v)$$

where

$$v(s) := Dg(\phi_\epsilon(\xi)(s))(D\phi_\epsilon(\xi)\xi'(s)), \quad s \leq 0.$$

Step 4: behaviour as $n \rightarrow \infty$.

Let $\xi_n \rightarrow \xi_0$ in $\mathbb{R}^\nu$ and let $\epsilon_n \rightarrow 0^+$. We claim that

$$(5.25) \quad \|\phi_{\epsilon_n}(\xi_n) - \phi_0(\xi_0)\|_{\zeta, \epsilon_n} \rightarrow 0 \quad \text{for } n \rightarrow \infty,$$

and for every $\xi' \in \mathbb{R}^\nu$

$$(5.26) \quad \|D\phi_{\epsilon_n}(\xi_n)(\xi') - D\phi_0(\xi_0)(\xi')\|_{\mu, \epsilon_n} \rightarrow 0 \quad \text{for } n \rightarrow \infty.$$

In fact,

$$\begin{aligned} \phi_{\epsilon_n}(\xi_n) - \phi_0(\xi_0) &= \Gamma_{\epsilon_n}(\xi_n, \phi_{\epsilon_n}(\xi_n)) - \Gamma_{\epsilon_n}(\xi_n, \phi_0(\xi_0)) \\ &\quad + \Gamma_{\epsilon_n}(\xi_n, \phi_0(\xi_0)) - \Gamma_0(\xi_0, \phi_0(\xi_0)). \end{aligned}$$

and

$$\|\Gamma_{\epsilon_n}(\xi_n, \phi_{\epsilon_n}(\xi_n)) - \Gamma_{\epsilon_n}(\xi_n, \phi_0(\xi_0))\|_{\zeta, \epsilon_n} \leq (1/2)\|\phi_{\epsilon_n}(\xi_n) - \phi_0(\xi_0)\|_{\zeta, \epsilon_n}$$

so

$$\begin{aligned} \|\phi_{\epsilon_n}(\xi_n) - \phi_0(\xi_0)\|_{\zeta, \epsilon_n} &\leq 2\|\Gamma_{\epsilon_n}(\xi_n, \phi_0(\xi_0)) - \Gamma_0(\xi_0, \phi_0(\xi_0))\|_{\zeta, \epsilon_n} \\ &= 2\|\Xi_{\epsilon_n, \nu}(\xi_n, y_n) - \Xi_{0, \nu}(\xi_0, y_0)\|_{\zeta, \epsilon_n} \\ &\leq 2(L + l)|\Xi_{\epsilon_n, \nu}(\xi_n, y_n) - \Xi_{0, \nu}(\xi_0, y_0)|_{\zeta, \epsilon_n} \end{aligned}$$

where $y_n \equiv y_0 = g \circ (\phi_0(\xi_0))$. Therefore (5.25) follows from Lemma 5.3. Now

$$(5.27) \quad |Dg(u)v|_{L^2} \leq L|v|_{L^2} + l|v|_{H^1} \leq \|v\|_\epsilon \quad \text{for } u, v \in H^1(\Omega).$$

Therefore, using (5.24) and proceeding as in the proof of (5.25), we obtain

$$\begin{aligned} &\|D\phi_{\epsilon_n}(\xi_n)(\xi') - D\phi_0(\xi_0)(\xi')\|_{\mu, \epsilon_n} \\ &\leq 2\|\Xi_{\epsilon_n, \nu}(\xi', y_n) - \Xi_{0, \nu}(\xi', y_0)\|_{\mu, \epsilon_n} \\ (5.28) \quad &\leq 2(L + l)|\Xi_{\epsilon_n, \nu}(\xi', y_n) - \Xi_{0, \nu}(\xi', y_0)|_{\mu, \epsilon_n} \end{aligned}$$

where, for every $s \leq 0$,

$$\begin{aligned} v(s) &:= D\phi_0(\xi_0)\xi'(s), \\ y_n(s) &:= Dg(\phi_{\epsilon_n}(\xi_n)(s))v(s) \end{aligned}$$

and

$$y_0(s) := Dg(\phi_0(\xi_0)(s))v(s).$$

We claim that

$$(5.29) \quad |y_n - y_0|_{\zeta, L^2} \quad \text{as } n \rightarrow \infty.$$

Assuming this claim, we obtain (5.26) from (5.28), (5.29) and Lemma 5.3. To prove the claim, note that by (5.27)

$$\begin{aligned} e^{\mu t} |y_n(t) - y_0(t)|_{L^2} &= e^{(\mu - \zeta)t} |Dg(\phi_{\epsilon_n}(\xi_n)(t))w(t) - Dg(\phi_0(\xi_0)(t))w(t)|_{L^2} \\ &\leq e^{(\mu - \zeta)t} \|v\|_{\eta, 0} \end{aligned}$$

where $w(t) := e^{\eta t} v(t)$, so $\|w(t)\|_{\epsilon_n} \|w(t)\|_0 \leq \|v\|_{\eta, 0}$, $t \leq 0$.

Therefore, given $\delta > 0$ there is a $a \leq 0$ such that

$$e^{\mu t} |y_n(t) - y_0(t)|_{\epsilon_n} \leq \delta \quad \text{for } t < a.$$

Thus we only need to prove that

$$\sup_{t \in [a, 0]} |Dg(\phi_{\epsilon_n}(\xi_n)(t))w(t) - Dg(\phi_0(\xi_0)(t))w(t)|_{L^2} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

However, this is obvious since $Dg: H^1(\Omega) \rightarrow L(H^1(\Omega), L^2(\Omega))$ is continuous and, by (5.25),

$$\sup_{t \in [a, 0]} |\phi_{\epsilon_n}(\xi_n)(t) - \phi_0(\xi_0)(t)|_{H^1} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Step 5: the global invariant manifold.

Define, for $0 < \epsilon \leq \epsilon_0$,

$$\Lambda_\epsilon: \mathbb{R}^\nu \rightarrow H^1(\Omega), \quad \xi \mapsto \phi_\epsilon(\xi)(0),$$

and

$$\Lambda_0: \mathbb{R}^\nu \rightarrow H_s^1(\Omega), \quad \xi \mapsto \phi_0(\xi)(0).$$

By what we have proved so far, for every $\epsilon \in [0, \epsilon_0]$ the map Λ_ϵ is well-defined, of class C^1 and (5.6) and (5.8) hold. It is well-known and easily proved that $\Lambda_\epsilon(\mathbb{R}^\nu)$ is an invariant manifold of the semiflow $\pi_{\epsilon, g}$ which includes all orbits of solutions of $\pi_{\epsilon, g}$ defined for $t \leq 0$ and lying in Z_ϵ^ζ . Since g equals $\hat{f}$ on U , it follows that every point in $\mathcal{A}_\epsilon$ is contained in $\Lambda_\epsilon(\mathbb{R}^\nu)$. The reduced equation on the manifold $\Lambda_\epsilon(\mathbb{R}^\nu)$ clearly takes the form (5.7) and (5.8) implies (5.9).

Step 6: the local invariant manifold.

Let

$$K := \{ \xi \in \mathbb{R}^\nu \mid V_0(\Lambda_0(\xi)) \leq M_0 \} = \{ \xi \in \mathbb{R}^\nu \mid \Lambda_0(\xi) \in B_2 \}.$$

Since B_2 is bounded and closed, it follows from (5.6) that K is bounded and closed, i.e. compact.

Define

$$V := \{ \xi \in \mathbb{R}^\nu \mid V_0(\Lambda_0(\xi)) < M_0 \}.$$

Thus $V \subset K$ and V is open in $\mathbb{R}^\nu$. Since $\Lambda(K) \subset U$ and K is compact and U is open in $H^1(\Omega)$, it follows from (5.8), by choosing $\epsilon_0 > 0$ smaller, if necessary, that

$$\Lambda_\epsilon(K) \subset U, \quad \epsilon \in [0, \epsilon_0].$$

We also claim that, if $\epsilon_0 > 0$ is small enough, then

$$\Lambda_\epsilon^{-1}(\mathcal{A}_\epsilon) \subset V, \quad \epsilon \in [0, \epsilon_0].$$

In fact if the claim does not hold, then there are sequences $\epsilon_n \rightarrow 0^+$ and $\xi_n \notin V$ with $u_n := \Lambda_{\epsilon_n}(\xi_n) \in \mathcal{A}_{\epsilon_n}$. By Corollary 5.2 and Lemma 5.9 of [15] we may assume that $u_n \rightarrow u_0$ in $H^1(\Omega)$, where $u_0 \in \mathcal{A}_0$. It follows that there is a $\xi_0 \in \mathbb{R}^\nu$ with $u_0 = \Lambda_0(\xi_0)$. Since, by (5.6), for every $a \in \mathbb{R}^\nu$ and every $\epsilon \in [0, \epsilon_0]$, the j -th component a_j of a is given by $a_j = \langle \Lambda_\epsilon(a), w_{\epsilon,j} \rangle$, it follows that $\xi_n \rightarrow \xi_0$ in $\mathbb{R}^\nu$. By the definition of V we have $\xi \in V$, a contradiction, proving the claim. Set $W := V_0 \circ \Lambda_0: \mathbb{R}^\nu \rightarrow \mathbb{R}$. Then, for every $\xi \in \Lambda_\epsilon^{-1}(U)$, we have

$$\begin{aligned} \nabla W(\xi) \cdot v_0(\xi) &= DV_0(\Lambda_0(\xi)) D\Lambda_0(\xi)(v_0(\xi)) \\ &= -|D\Lambda_0(\xi)(v_0(\xi))|_{L^2}^2. \end{aligned}$$

Since there are no equilibria u of $\pi_{0,\hat{f}}$ with $V_0(u) = M_0$, it follows that

$$D\Lambda_0(\xi)(v_0(\xi)) \neq 0, \quad \text{whenever } W(\xi) = M_0.$$

By the compactness of K we now obtain that there is a $\delta > 0$ such that

$$\nabla W(\xi) \cdot v_0(\xi) < -\delta, \quad \text{whenever } W(\xi) = M_0.$$

Therefore (5.9) implies that, if $\epsilon_0 > 0$ is small enough, then

$$\nabla W(\xi) \cdot v_\epsilon(\xi) < -\delta, \quad \text{whenever } W(\xi) = M_0.$$

This shows, that, for $\epsilon \in [0, \epsilon_0]$, the set V is positively invariant for the equation (5.7) so the set $\Lambda_\epsilon(V) \subset U$ is positively invariant for the semiflow $\pi_{\epsilon,\hat{f}}$ and $\mathcal{A}_\epsilon \subset \Lambda_\epsilon(V)$. The theorem is proved. $\square$

6. FINAL REMARKS

It is clear that Theorem 5.2, by its abstract nature, can easily be generalized to the case of an arbitrary smooth bounded domain $\Omega \subset \mathbb{R}^M \times \mathbb{R}^N$, with $n := M + N \geq 2$. In this case the squeezed domain Ω_ϵ is defined by

$$\Omega_\epsilon := \{ (x, \epsilon y) \mid (x, y) \in \Omega \},$$

where now $x \in \mathbb{R}^M$ and $y \in \mathbb{R}^N$. If $n > 2$ one only needs to impose the condition

$$(6.1) \quad \beta \leq (2^*/2) - 1, \quad \text{where } 2^* = 2n/(n-2)$$

in (2.1), so the Nemitski operator associated with f is a C^1 -map of H^1 into L^2 (see [15] for details). The strict inequality in (6.1) makes it possible to use the Gagliardo-Nirenberg inequality in the proof of Theorem 5.1.




Figure 3

Of course, this generalization is meaningful only if one knows that the eigenvalues of the limit operator A_0 satisfy the gap condition (5.5). Under fairly general conditions, this is actually the case if $M = 1$ and $N \geq 1$ is arbitrary. In fact, one can easily extend the notion of *nicely decomposed domain* in order to cover this more general situation. If the additional condition (C) introduced in Section 4 is satisfied, then one can prove (5.5) by just repeating the arguments of Section 4. We do not enter into detail: Figure 3 illustrates a simple example of a three-dimensional domain on which this kind of analysis is possible.

If $M \geq 2$ the problem of the spectral gap condition is much more delicate (see [12] and [13] for this and some related questions). Some (partial) generalizations of Theorem 5.2 should be possible for very particular three-dimensional domains, essentially for domains which are ordinate sets of positive functions defined on rectangles.

One can also consider, much more generally, a squeezing transformation of an open subset of $\mathbb{R}^{M+N}$ toward an arbitrary smooth M -dimensional submanifold S of $\mathbb{R}^{M+N}$ (see [14] for details). If the eigenvalues of the Laplace-Beltrami operator on S satisfy a gap condition like (5.5) (e.g. when S is a sphere), some further generalizations of Theorem 5.2 have been obtained in [16].

7. APPENDIX

In this appendix we give the

Proof of Lemma 4.4. First of all, we observe that, since $q''(x) \leq 0$ on $]0, 1]$, the function q is concave on $]0, 1[$. So, for $x \in]0, 1[$ and for all $0 < \epsilon < x$, we have $q(x) + (\epsilon - x)q'(x) \geq q(\epsilon)$. Now, since q is continuous on $[0, 1]$ and $q(0) = 0$, we get

$$(7.1) \quad 0 \leq (q'(x)/q(x)) \leq (1/x) \quad \text{for } x \in]0, 1].$$

Next, we analyze some properties of eigenvectors of (a^q, b^q) . Let λ be an eigenvalue of (a^q, b^q) , and let $u \in V(q)$ be a corresponding eigenvector. This means that

$$\int_0^1 qu'v' dx = \lambda \int_0^1 quv dx \quad \text{for all } v \in V(q).$$

Choosing $v \in C_0^\infty(]0, 1])$ arbitrarily, we see that $(qu')' \in L_{\text{loc}}^1(0, 1)$ and $(qu')' = -\lambda qu$. In particular, we see that $(qu')' \in C^0([0, 1])$, and hence $qu' \in C^1([0, 1])$. Moreover, taking v any function in $C^\infty([0, 1])$, with $v \equiv 0$ on $[0, \bar{\epsilon}]$, $v \equiv 1$ on $[1 - \bar{\epsilon}, 1]$, where $0 < \bar{\epsilon} < 1 - \bar{\epsilon}$, we obtain for any ϵ , $0 < \epsilon < \bar{\epsilon}$:

$$\begin{aligned} \lambda \int_0^1 quv dx &= \int_0^1 qu'v' dx = \int_\epsilon^{1-\epsilon} qu'v' dx \\ &= - \int_\epsilon^{1-\epsilon} (qu')'v dx + q(1-\epsilon)u'(1-\epsilon)v(1-\epsilon) - q(\epsilon)u'(\epsilon)v(\epsilon) \\ &= \lambda \int_\epsilon^{1-\epsilon} quv dx + q(1-\epsilon)u'(1-\epsilon). \end{aligned}$$

By letting $\epsilon \rightarrow 0$, we obtain that $u'(1) = 0$. Taking now v any function in $C^\infty([0, 1])$, with $v \equiv 1$ in $[0, \bar{\epsilon}]$, $v \equiv 0$ on $[1 - \bar{\epsilon}, 1]$, where $0 < \bar{\epsilon} < 1 - \bar{\epsilon}$, we obtain for any ϵ , $0 < \epsilon < \bar{\epsilon}$:

$$\begin{aligned} \lambda \int_0^1 quv dx &= \int_0^1 qu'v' dx = \int_\epsilon^{1-\epsilon} qu'v' dx \\ &= - \int_\epsilon^{1-\epsilon} (qu')'v dx + q(1-\epsilon)u'(1-\epsilon)v(1-\epsilon) - q(\epsilon)u'(\epsilon)v(\epsilon) \\ &= \lambda \int_\epsilon^{1-\epsilon} quv dx - q(\epsilon)u'(\epsilon). \end{aligned}$$

By letting $\epsilon \rightarrow 0$, we obtain that $(qu')(0) = 0$. But now, for $x \in]0, 1]$, we have:

$$(qu')(x) = \int_0^x (qu')'(s) ds = -\lambda \int_0^x q(s)u(s) ds;$$

since q is nonnegative and nondecreasing on $[0, 1]$, it follows that

$$|q(x)u'(x)| \leq \lambda \int_0^x q(s)|u(s)| ds \leq \lambda q(x)|u|_\infty x,$$

and, since $q(x) > 0$ for $x \in]0, 1]$,

$$(7.2) \quad |u'(x)| \leq \lambda |u|_\infty x.$$

In particular, $u \in C^1([0, 1])$ and $u'(0) = 0$.

Assume now that $\lambda \neq 0$. Then $u' \not\equiv 0$. Let us define

$$w(x) := q(x)^{1/2} u'(x), \quad x \in [0, 1].$$

Then $w \in C^0([0, 1])$ and $w(0) = w(1) = 0$. Moreover, in $]0, 1]$ we have:

$$\begin{aligned} w' &= (q^{-1/2} q u')' = -\frac{1}{2} q^{-3/2} q' q u' + q^{-1/2} (q u')' \\ &= -\frac{1}{2} \frac{q'}{q} q^{1/2} u' - q^{-1/2} \lambda q u \\ &= -\frac{1}{2} \frac{q'}{q} w - \lambda q^{1/2} u. \end{aligned}$$

From this equality, it follows that $w' \in C^0([0, 1])$, so $w \in C^1([0, 1])$; but then the same equality implies that $w' \in C^1([0, 1])$, so $w \in C^2([0, 1])$. Moreover, (7.1) and (7.2) imply that $w' \in L^2(0, 1)$, so $w \in H_0^1$. We compute w'' :

$$\begin{aligned} w'' &= -\frac{1}{2} \left(\frac{q'}{q} \right)' w - \frac{1}{2} \frac{q'}{q} w' - \lambda (q^{1/2} u)' \\ &= -\frac{1}{2} \left(\frac{q'}{q} \right)' w - \frac{1}{2} \frac{q'}{q} \left(-\frac{1}{2} \frac{q'}{q} w - \lambda q^{1/2} u \right) - \frac{1}{2} \lambda q^{-1/2} q' u - \lambda w \\ &= -\frac{1}{2} \left(\frac{q'}{q} \right)' w + \frac{1}{4} \frac{q'^2}{q^2} w - \lambda w \\ &= -\frac{1}{2} \frac{q'' q - q'^2}{q^2} w + \frac{1}{4} \frac{q'^2}{q^2} w - \lambda w \\ &= \left(\frac{3}{4} \frac{q'^2}{q^2} - \frac{1}{2} \frac{q''}{q} \right) w - \lambda w \end{aligned}$$

Let us define

$$Q(x) := \frac{3}{4} \frac{q'(x)^2}{q(x)^2} - \frac{1}{2} \frac{q''(x)}{q(x)}, \quad x \in]0, 1].$$

Observe that $0 \leq Q(x) \leq -(3/4)(q'/q)'(x)$ on $]0, 1]$. We claim that $Q^{1/2} w \in$

$L^2(0, 1)$. In fact let ϵ , $0 < \epsilon < 1$, be arbitrary: by (7.1) and (7.2), we have:

$$\begin{aligned}
\int_{\epsilon}^1 Q w^2 dx &\leq \frac{3}{4} \int_{\epsilon}^1 \left(-\frac{q'}{q} \right)' w^2 dx \\
&\leq \frac{3}{4} \int_{\epsilon}^1 \left(-\frac{q'}{q} \right)' (\lambda |u|_{\infty} |q|_{\infty}^{1/2} x)^2 dx \\
&\leq \frac{3}{4} \lambda |u|_{\infty}^2 |q|_{\infty} \int_{\epsilon}^1 \left(-\frac{q'}{q} \right)' x^2 dx \\
&= \frac{3}{4} \lambda |u|_{\infty}^2 |q|_{\infty} \left(-\frac{q'}{q} x^2 \Big|_{\epsilon}^1 + 2 \int_{\epsilon}^1 \frac{q'}{q} x dx \right) \\
&\leq \frac{3}{4} \lambda |u|_{\infty}^2 |q|_{\infty} \left(\frac{1}{\epsilon} \epsilon^2 + 2 \int_{\epsilon}^1 \frac{1}{x} dx \right).
\end{aligned}$$

Since ϵ is arbitrary, the claim is proved.

Let us define the Hilbert space

$$U(Q) := \left\{ \omega \in H_0^1(0, 1) \mid Q^{1/2} \omega \in L^2(0, 1) \right\},$$

with the scalar product

$$\langle \omega, \varpi \rangle_{U(Q)} := \int_0^1 \omega' \varpi' dx + \int_0^1 Q \omega \varpi dx.$$

Let us notice that $U(Q) \hookrightarrow H_0^1(0, 1)$ with continuous imbedding, so the imbedding $U(Q) \hookrightarrow L^2(0, 1)$ is compact. Moreover, $U(Q)$ is clearly a dense subspace of $L^2(0, 1)$, since it contains $C_0^{\infty}(]0, 1[)$. The hypothesis of Proposition 2.2 in [15] is then satisfied by the pair (a_Q, b) , where $a_Q := \langle \cdot, \cdot \rangle_{U(Q)}$ and $b := \langle \cdot, \cdot \rangle_{L^2(0, 1)}$. Then it follows that there exists an eigenvalue-eigenvector sequence $(\mu_{\nu}^Q, \omega_{\nu}^Q)_{\nu \in \mathbb{N}}$ of the pair (a_Q, b) such that

$$0 \leq \mu_1^Q \leq \mu_2^Q \leq \mu_3^Q \leq \dots$$

and $(\omega_{\nu}^Q)_{\nu \in \mathbb{N}}$ is a complete orthonormal system in $L^2(0, 1)$. We have proved above that $w \in U(Q)$. We claim that

$$\int_0^1 w' \omega' dx + \int_0^1 Q w \omega dx = \lambda \int_0^1 w \omega dx \quad \text{for all } \omega \in U(Q),$$

i.e. (λ, w) is an eigenvalue-eigenvector pair of (a_Q, b) . In fact, for every ϵ , $0 < \epsilon < 1$, we have

$$\begin{aligned}
\int_{\epsilon}^1 w' \omega' dx &= - \int_{\epsilon}^1 w'' \omega dx + w' \omega|_{\epsilon}^1 \\
&= - \int_{\epsilon}^1 Q w \omega dx + \lambda \int_{\epsilon}^1 w \omega dx - w'(\epsilon) \omega(\epsilon).
\end{aligned}$$

Then

$$\int_0^1 w' \omega' dx + \int_0^1 Q w \omega dx = \lambda \int_0^1 w \omega dx - \lim_{\epsilon \rightarrow 0} w'(\epsilon) \omega(\epsilon).$$

Recall that

$$w'(\epsilon) = \left(-\frac{1}{2} \frac{q'(\epsilon)}{q(\epsilon)} q(\epsilon)^{1/2} u'(\epsilon) - \lambda q(\epsilon)^{1/2} u(\epsilon) \right);$$

so, by (7.1) and (7.2), it follows that $w'(\epsilon) \omega(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$ and the claim is proved.

Thus we have proved that, whenever (λ, u) is an eigenvalue-eigenvector pair for (a^q, b^q) and $\lambda \neq 0$, then $(\lambda, q^{1/2} u')$ is an eigenvalue-eigenvector pair for (a_Q, b) . Assume now that $\lambda \neq 0$ is an eigenvalue of (a^q, b^q) , and that $u, \tilde{u} \in V(q)$ are two linearly independent corresponding eigenvectors. Let $w := q^{1/2} u'$ and $\tilde{w} := q^{1/2} \tilde{u}'$. Then also w and $\tilde{w}$ are linearly independent. Otherwise, we could find $\xi \neq 0$ such that $w = \xi \tilde{w}$, that is $q^{1/2} u' = \xi q^{1/2} \tilde{u}'$. Since $q(x) > 0$ on $]0, 1]$, it follows that $u' = \xi \tilde{u}'$. So there is a constant ζ such that $u = \xi \tilde{u} + \zeta$, that is $u, \tilde{u}$ and 1 are linearly dependent, a contradiction. This means that the multiplicity of λ as an eigenvalue of (a^q, b^q) is less than or equal to the multiplicity of λ as an eigenvalue of (a_Q, b) . So we can conclude that

$$\lambda_1^q = 0 \quad \text{and} \quad \lambda_\nu^q \geq \mu_{\nu-1}^Q \quad \text{for } \nu \geq 2.$$

In order to complete the proof, we need to show that $\mu_\nu^Q \geq \pi^2 \nu^2$, for $\nu = 1, 2, \dots$. This is done as follows: by Proposition 2.2 in [15],

$$\mu_\nu^Q = \inf_{E \in \mathcal{U}_\nu(Q)} \sup_{\omega \in E \setminus \{0\}} \frac{\int_0^1 \omega'^2 dx + \int_0^1 Q \omega^2 dx}{\int_0^1 \omega^2 dx},$$

where $\mathcal{U}_\nu(Q)$ is the set of all ν -dimensional subspaces of $U(Q)$. Since $Q \geq 0$ and $U(Q) \subset H_0^1(0, 1)$, it follows that

$$\mu_\nu^Q \geq \mu_\nu^0 := \inf_{E \in \mathcal{U}_\nu} \sup_{\omega \in E \setminus \{0\}} \frac{\int_0^1 \omega'^2 dx}{\int_0^1 \omega^2 dx},$$

where $\mathcal{U}_\nu$ is the set of all ν -dimensional subspaces of $H_0^1(0, 1)$. But $(\mu_\nu^0)_{\nu \in \mathbb{N}}$ are exactly the eigenvalues of the Dirichlet problem

$$\begin{cases} \omega''(x) = \mu \omega(x), & x \in]0, 1[\\ u(x) = 0, & x \in \{0, 1\} \end{cases}$$

This implies that $\mu_\nu^0 = \pi^2 \nu^2$, $\nu = 1, 2, \dots$, and the proof is complete. $\square$

REFERENCES

1. A. Ambrosetti and G. Prodi, *Analisi non lineare*, I Quaderno, Scuola Normale Superiore – Classe di Scienze, Pisa, 1973.
2. J. Arrieta, *Spectral behavior and upper semicontinuity of attractors*, International Conference on Differential Equations vol. 1,2 (Berlin, 1999), World Sci. Publishing, River Edge, NJ (2000), 615–621.
3. P. Brunovský and I. Tereščák, *Regularity of invariant manifolds*, J. Dyn. Differ. Equations **3** (1991), 313 – 337.
4. S. N. Chow and K. Lu, *Invariant manifolds for flows in Banach spaces*, J. Differ. Equations **74** (1988), 285 – 317.
5. I.S. Ciuperca, *Spectral properties of Schrödinger operators on domains with varying order of thinness*, J. Dyn. Differ. Equations **10** (1998), 73 – 108.
6. Jack K. Hale, *Asymptotic Behavior of Dissipative Systems*, Math. Surveys Monographs 25, AMS, Providence, 1988.
7. J. Hale and G. Raugel, *Reaction-diffusion equations on thin domains*, J. Math. Pures Appl. **IX Ser.71** (1992), 33 – 95.
8. J. Hale and G. Raugel, *A damped hyperbolic equation on thin domains*, Trans. Am. Math. Soc. **329** (1992), 185 – 219.
9. J. Hale and G. Raugel, *A reaction-diffusion equation on a thin L-shaped domain*, Proc. Roy. Soc. Edinb., Sect A **125** (1995), 283 – 327.
10. D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Lecture notes in mathematics, Vol 840, Springer-Verlag, NY, 1981.
11. O. Ladyzhenskaya, *Attractors for Semigroups and Evolution Equations*, Cambridge University Press, Cambridge, 1991.
12. J. Mallet-Paret and G.R. Sell, *Inertial manifolds for reaction-diffusion equations in higher space dimensions*, Jour. Am. Math. Society **1** (1988), 805 – 866.
13. J. Mallet-Paret, G.R. Sell and Z.D. Shao, *Obstructions to the existence of normally hyperbolic inertial manifolds*, Indiana Univ. Math. J. **42** (1993), 1027–1055.
14. M. Prizzi, M. Rinaldi and K. P. Rybakowski, *Curvilinear domain squeezing and parabolic equations*, Studia Math. **151** (2002), 109–140.
15. M. Prizzi and K. P. Rybakowski, *The effect of domain squeezing upon the dynamics of reaction-diffusion equations*, J. Differ. Equations **173** (2001), 271 – 320.
16. M. Prizzi and K. P. Rybakowski, *On inertial manifolds for reaction-diffusion equations on genuinely high-dimensional thin domains*, Studia Math. (to appear).
17. G. Raugel, *Dynamics of partial differential equations on thin domains*, R. Johnson (ed.), Dynamical systems. Lectures given at the 2nd session of the Centro Internazionale Matematico Estivo (CIME) held in Montecatini Terme, Italy, June 13 – 22, 1994. Lecture notes in mathematics, Vol. 1609, Springer-Verlag, Berlin, 1995, pp. 208 – 315.
18. K. P. Rybakowski, *An abstract approach to smoothness of invariant manifolds*, Appl. Analysis **49** (1993), 119 – 150.

UNIVERSITÀ DEGLI STUDI DI TRIESTE, DIPARTIMENTO DI SCIENZE MATEMATICHE, VIA VALERIO 12/B, 34100 TRIESTE, ITALY

E-mail address: prizzi@mathsun1.univ.trieste.it

UNIVERSITÄT ROSTOCK, FACHBEREICH MATHEMATIK, UNIVERSITÄTSPLATZ 1, 18055 ROSTOCK, GERMANY

E-mail address: krzysztof.rybakowski@mathematik.uni-rostock.de